arXiv:1502.07415v1 [math.RT] 26 Feb 2015

SYMMETRIC QUIVER HECKE ALGEBRAS AND R-MATRICES OF QUANTUM AFFINE ALGEBRAS IV

SEOK-JIN KANG[1], MASAKI KASHIWARA[2], MYUNGHO KIM AND SE-JIN OH[3]

Abstract. Let $U'_q(\mathfrak{g})$ be a twisted affine quantum group of type $A_N^{(2)}$ or $D_N^{(2)}$ and let $\mathfrak{g}_0$ be the finite-dimensional simple Lie algebra of type A_N or D_N. For a Dynkin quiver of type $\mathfrak{g}_0$, we define a full subcategory $\mathcal{C}_Q^{(2)}$ of the category of finite-dimensional integrable $U'_q(\mathfrak{g})$-modules, a twisted version of the category $\mathcal{C}_Q$ introduced by Hernandez and Leclerc. Applying the general scheme of affine Schur-Weyl duality, we construct an exact faithful KLR-type duality functor $\mathcal{F}_Q^{(2)}\colon \mathrm{Rep}(R) \to \mathcal{C}_Q^{(2)}$, where $\mathrm{Rep}(R)$ is the category of finite-dimensional modules over the quiver Hecke algebra R of type $\mathfrak{g}_0$ with nilpotent actions of the generators x_k. We show that $\mathcal{F}_Q^{(2)}$ sends any simple object to a simple object and induces a ring isomorphism $\mathrm{K}(\mathrm{Rep}(R)) \xrightarrow{\sim} \mathrm{K}(\mathcal{C}_Q^{(2)})$.

Introduction

This paper is a continuation of our previous works on symmetric quiver Hecke algebras and R-matrices of quantum affine algebras [15, 16, 18]. In particular, this work can be considered as a twisted version of [16].

The classical Schur-Weyl duality explains the close connection between the representation theory of the symmetric group $\mathfrak{S}_k$ and the representation theory of general linear Lie algebra $\mathfrak{gl}_n$ [31, 32]. In [13], Jimbo introduced a quantum version of Schur-Weyl duality between the Hecke algebra H_k and quantized universal enveloping algebra $U_q(\mathfrak{gl}_n)$. In [6, 7, 9], Chari-Pressley, Cherednik, Ginzburg-Reshetikhin-Vasserot constructed a quantum affine Schur-Weyl duality functor from the category of finite-dimensional representations of the affine Hecke algebra H_k^{aff} to the category of finite-dimensional integrable modules over the quantum affine algebra $U_q(A_N^{(1)})$.

Date: February 25, 2015.

[1] This work was supported by NRF Grant # 2014-021261 and by NRF Grant # 2010-0010753.

[2] This work was partially supported by Grant-in-Aid for Scientific Research (B) 22340005, Japan Society for the Promotion of Science.

[3] This work was supported by BK21 PLUS SNU Mathematical Sciences Division.

Using the symmetric quiver Hecke algebras and R-matrices of quantum affine algebras, it was shown in [15] that one can construct quantum affine Schur-Weyl duality functors in a more general setting. Let $\{V_s\}_{s\in\mathcal{S}}$ be a family of good modules over a quantum affine algebra $U_q'(\mathfrak{g})$. By investigating the distribution of poles of normalized R-matrices between the V_s's, we obtain a symmetric quiver Hecke algebra R and a $(U_q'(\mathfrak{g}),R)$-bimodule $\widehat{V}$. In this way, we construct the *KLR-type quantum affine Schur-Weyl duality functor* (or *KLR-type duality functor* for brevity) $\mathcal{F}$ from the category $\mathrm{Rep}(R)$ to the category $\mathcal{C}_\mathfrak{g}$ of finite-dimensional integrable $U_q'(\mathfrak{g})$-modules given by $M\mapsto \widehat{V}\otimes_R M$ $(M\in\mathrm{Rep}(R))$. Here, $\mathrm{Rep}(R)$ is the category of finite-dimensional modules over the quiver Hecke algebra R on which the generators x_k act nilpotently. When R is of finite A, D, E type, $\mathcal{F}$ is exact ([23]). With various choices of quantum affine algebras and good modules, one can construct a lot of interesting and significant KLR-type duality functors.

In [15, 18], by taking $\{V(\varpi_1)_{(-q)^{2s}}\}_{s\in\mathbb{Z}}$ as the family of good modules over $U_q(A_N^{(t)})$ $(N\in\mathbb{Z}_{\ge 2},\ t\in\{1,2\})$, we constructed the exact functor $\mathcal{F}$ from $\mathrm{Rep}(R)$ to $\mathcal{C}^0_{A_N^{(t)}}$, and investigated properties of $\mathcal{F}$ and the relationships among the categories $\mathrm{Rep}(R)$, $\mathcal{C}^0_{A_N^{(1)}}$ and $\mathcal{C}^0_{A_N^{(2)}}$. Here $\mathcal{C}^0_\mathfrak{g}$ is the category of finite-dimensional modules over the quantum affine algebra $U_q'(\mathfrak{g})$ introduced in [11] and R is the quiver Hecke algebra of type A_∞.

In [16], motivated by the work of Hernandez and Leclerc [12], Kang, Kashiwara and Kim applied the general scheme to the quantum affine algebras $U_q'(A_N^{(1)})$, $U_q'(D_N^{(1)})$ and their fundamental representations. Let $\mathfrak{g}^{(1)}$ be an untwisted affine Kac-Moody algebra of type $A_N^{(1)}$ or $D_N^{(1)}$ and let $\mathfrak{g}_0^{(1)}$ be the classical part of $\mathfrak{g}^{(1)}$, the finite-dimensional simple Lie algebra of type A_N or D_N inside $\mathfrak{g}^{(1)}$. We denote by $I^{(1)}$ and $I_0^{(1)}$ the index set of simple roots for $\mathfrak{g}^{(1)}$ and $\mathfrak{g}_0^{(1)}$, respectively. Let $\Delta^+_{\mathfrak{g}_0^{(1)}}$ be the set of positive roots of $\mathfrak{g}_0^{(1)}$ and $\Pi_{\mathfrak{g}_0^{(1)}}$ the set of simple roots. For any Dynkin quiver Q of type $\mathfrak{g}_0^{(1)}$, we set $\Gamma_Q=\phi^{-1}(\Delta^+_{\mathfrak{g}_0^{(1)}}\times\{0\})\subset I_0^{(1)}\times\mathbb{Z}$, the Auslander-Reiten quiver of Q. (See Section 2 for more details including the definition of ϕ.) We take the fundamental representations $V(\varpi_i)$ $(i\in I_0^{(1)})$ as the family of good modules over $U_q'(\mathfrak{g}^{(1)})$ and take the index set $J=\phi^{-1}(\Pi_{\mathfrak{g}_0^{(1)}}\times\{0\})$ together with the maps $s^{(1)}\colon J\to I_0^{(1)}$ and $X^{(1)}\colon J\to\mathbf{k}^\times$ given by $s^{(1)}\colon (i,p)\mapsto i$, $X^{(1)}\colon (i,p)\mapsto(-q)^p$ for $(i,p)\in J$. Then the corresponding quiver Hecke algebra R is of type $\mathfrak{g}_0^{(1)}$ (i.e., A_N or D_N) and the KLR-type duality functor is an exact functor $\mathcal{F}_Q^{(1)}\colon\mathrm{Rep}(R)\to\mathcal{C}_Q^{(1)}$, where $\mathcal{C}_Q^{(1)}$ is the full subcategory of $\mathcal{C}_{\mathfrak{g}^{(1)}}$ introduced in [12].

Moreover, $\mathcal{F}_Q^{(1)}$ sends any simple object to a simple object and induces a ring isomorphism $\mathrm{K}(\mathrm{Rep}(R)) \xrightarrow{\sim} \mathrm{K}(\mathcal{C}_Q^{(1)})$.

In this paper, we extend these results on $\mathfrak{g}^{(1)}$ to the twisted quantum affine algebras $U_q'(\mathfrak{g})$ of type $A_N^{(2)}$ $(N \ge 2)$ and $D_N^{(2)}$ $(N \ge 4)$ (see Remark 3.1). As in the untwisted case, we take the fundamental representations $V(\varpi_i)$ $(i \in I_0)$ as the family of good modules. But to take care of twisted setting, we need to develop some *twisted* techniques.

Let $\mathfrak{g}^{(t)} = A_N^{(t)}$ $(t = 1, 2)$ or $\mathfrak{g}^{(t)} = D_N^{(t)}$ $(t = 1, 2)$. Let $\mathfrak{g}_0^{(t)}$ be the classical part of $\mathfrak{g}^{(t)}$. Let $I^{(t)}$ and $I_0^{(t)}$ be the index set of simple roots for $\mathfrak{g}^{(t)}$ and $\mathfrak{g}_0^{(t)}$, respectively.

For an arbitrary affine Kac-Moody algebra $\mathfrak{g}$, we denote by $\mathscr{S}(\mathfrak{g})$ the quiver with the evaluation modules of fundamental modules of $U_q'(\mathfrak{g})$ as vertices and with the arrows determined by the poles of the normalized R-matrices (see (3.5)).

Then we have a quiver isomorphism

$$\pi_{\mathfrak{g}^{(1)}} \colon \mathscr{S}_0(\mathfrak{g}^{(1)}) \xrightarrow{\sim} \mathscr{S}_0(\mathfrak{g}^{(2)}). \tag{0.1}$$

where $\mathscr{S}_0(\mathfrak{g}^{(t)})$ is a connected component of $\mathscr{S}(\mathfrak{g}^{(t)})$ $(t = 1, 2)$ (Proposition 3.3).

Set $J = \phi^{-1}(\Pi_{\mathfrak{g}_0^{(1)}} \times \{0\})$. Then we have defined a map $J \to I_0^{(1)} \times \mathbf{k}^\times \simeq \mathscr{S}(\mathfrak{g}^{(1)})$. Its image is contained in $\mathscr{S}_0(\mathfrak{g}^{(1)})$. Composing it with (0.1) we obtain $s^{(2)} \colon J \to I_0^{(2)}$ and $X^{(2)} \colon J \to \mathbf{k}^\times$. Then the quiver $\mathscr{S}^J$ associated with J and $(X^{(2)}, s^{(2)})$ (see (2.2)) coincides with Q^{rev}, the reversed quiver of Q. Hence the associated quiver Hecke algebra R is of type $\mathfrak{g}_0^{(1)}$, i.e. $R = R^{A_N}$ or R^{D_N}. Thus, we obtain an exact KLR-type duality functor

$$\mathcal{F}_Q^{(2)} \colon \mathrm{Rep}(R) \to \mathcal{C}_{\mathfrak{g}^{(2)}}.$$

Note that we have already constructed an exact functor $\mathcal{F}_Q^{(1)} \colon \mathrm{Rep}(R) \to \mathcal{C}_{\mathfrak{g}^{(1)}}$ with the same quiver Hecke algebra R.

The main results of this paper are the following:

(i) $\mathcal{F}_Q^{(2)}$ sends the simple objects to simple objects.

(ii) $\mathcal{F}_Q^{(2)}$ gives a bijection between the set of simple objects in $\mathrm{Rep}(R)$ and the set of simple objects in the full subcategory $\mathcal{C}_Q^{(2)}$ of $\mathcal{C}_{\mathfrak{g}^{(2)}}$ defined in Definition 4.3.

(iii) $\mathcal{F}_Q^{(2)}$ restricts to a faithful functor from $\mathrm{Rep}(R)$ to the category $\mathcal{C}_Q^{(2)}$.

(iv) $\mathcal{F}_Q^{(2)}$ induces a ring isomorphism $\mathrm{K}(\mathrm{Rep}(R)) \xrightarrow{\sim} \mathrm{K}(\mathcal{C}_Q^{(2)})$.

(v) The ring isomorphism $\mathrm{K}(\mathcal{C}_Q^{(1)}) \xrightarrow{\sim} \mathrm{K}(\mathcal{C}_Q^{(2)})$, which is obtained by (iv) and the ring isomorphism $\mathrm{K}(\mathrm{Rep}(R)) \xrightarrow{\sim} \mathrm{K}(\mathcal{C}_Q^{(1)})$, preserves the dimension.

Note that a similar relationship between $\mathcal{C}_{\mathfrak{g}^{(1)}}$ and $\mathcal{C}_{\mathfrak{g}^{(2)}}$ have been also studied by Hernandez [10] with an approach using q-character. He showed that there is a ring isomorphism between the Grothendieck rings $K(\mathcal{C}_{\mathfrak{g}^{(1)}})$ and $K(\mathcal{C}_{\mathfrak{g}^{(2)}})$, which sends Kirillov-Reshetikhin modules to Kirillov-Reshetikhin modules. This is one of our motivations.

This paper is organized as follows. In Section 1, we briefly review some of basic knowledge on quiver Hecke algebras and quantum affine algebras. In Section 2, we recall the general construction of KLR-type duality functors and its application to untwisted quantum affine algebras. In Section 3, we investigate the relation between untwisted and twisted quantum affine algebras. In Section 4, we prove the main results of our paper. Section 5 deals with the existence of non-zero $U_q'(\mathfrak{g})$-module homomorphisms between evaluation modules of fundamental representations and their tensor products.

1. Quiver Hecke algebras and quantum affine algebras

1.1. Quantum groups associated with Cartan datum. In this subsection, we recall the definition of quantum groups. Let I be an index set. A Cartan datum $(\mathsf{A}, \mathsf{P}, \mathsf{P}^\vee, \Pi, \Pi^\vee)$ consists of

(1) a generalized Cartan matrix $\mathsf{A} = (a_{ij})_{i,j\in I}$ satisfying

$$a_{ii} = 2 \ (i \in I), \quad a_{ij} \in \mathbb{Z}_{\le 0} \ (i \ne j) \quad \text{and} \quad a_{ij} = 0 \iff a_{ji} = 0,$$

(2) a free abelian group P, called the *weight lattice*,
(3) $\mathsf{P}^\vee = \mathrm{Hom}(\mathsf{P}, \mathbb{Z})$, called the *co-weight lattice*,
(4) $\Pi = \{\alpha_i \mid i \in I\}$, called the *set of simple roots*,
(5) $\Pi^\vee = \{h_i \mid i \in I\}$, called the *set of simple co-roots*,

such that

(a) $\langle h_i, \alpha_j\rangle = a_{ij}$ for all $ij \in I$,
(b) Π and $\Pi^\vee$ are linearly independent sets,
(c) for every $i \in I$, there exists $\Lambda_i \in \mathsf{P}$ such that $\langle h_j, \Lambda_i\rangle = \delta_{i,j}$.

We say that A is *symmetrizable* if there exists a diagonal matrix $\mathsf{D} = \mathrm{diag}(\mathsf{s}_i \in \mathbb{Z}_{>0} \mid i \in I)$ such that DA is symmetric. We call $\mathsf{Q} := \bigoplus_{i\in I} \mathbb{Z}\alpha_i$ the *root lattice* and $\mathsf{Q}^+ := \sum_{i\in I} \mathbb{Z}_{\ge 0}\alpha_i$ the *positive root lattice*. For $\beta = \sum_{i\in I} n_i\alpha_i \in \mathsf{Q}^+$, we set

$$\mathrm{ht}(\beta) = \sum_{i\in I} n_i \in \mathbb{Z}_{\ge 0} \quad \text{and} \quad \mathrm{supp}(\beta) = \{i \in I \mid n_i > 0\}.$$

Let A be a symmetrizable generalized Cartan matrix and let q be an indeterminate. Let $(\ ,\ )$ be a non-degenerate symmetric bilinear form on $\mathbb{Q}\otimes_{\mathbb{Z}}\mathsf{P}$ satisfying $\langle h_i,\lambda\rangle = \dfrac{2(\alpha_i,\lambda)}{(\alpha_i,\alpha_i)}$ for every $i\in I$ and $\lambda\in\mathbb{Q}\otimes_{\mathbb{Z}}\mathsf{P}$.

For $i\in I$ and $m\in\mathbb{Z}$, we set $q_i := q^{\frac{(\alpha_i,\alpha_i)}{2}}$, $[m]_i = (q_i^m - q_i^{-m})/(q_i - q_i^{-1})$ and $[m]_i! = \prod_{k=1}^m [k]_i$. Let us denote by d the smallest positive integer such that $d\frac{(\alpha_i,\alpha_i)}{2}\in\mathbb{Z}$ for all $i\in I$.

Definition 1.1. The *quantum group* $U_q(\mathfrak{g})$ associated with a Cartan datum $(\mathsf{A},\mathsf{P},\mathsf{P}^\vee,\Pi,\Pi^\vee)$ is the associative $\mathbb{Q}(q^{1/d})$-algebra generated by e_i, f_i $(i\in I)$ and q^h $(h\in d^{-1}\mathsf{P}^\vee)$ subject to the following relations:

(i) $q^0=1$, $q^hq^{h'}=q^{h+h'}$ for $h,h'\in d^{-1}\mathsf{P}^\vee$,

(ii) $q^he_i = q^{\langle h,\alpha_i\rangle}e_iq^h$, $q^hf_i = q^{-\langle h,\alpha_i\rangle}f_iq^h$ for $h\in d^{-1}\mathsf{P}^\vee$,

(iii) $e_if_j - f_je_i = \dfrac{K_i-K_i^{-1}}{q_i-q_i^{-1}}$ for $K_i := q^{\frac{(\alpha_i,\alpha_i)}{2}h_i}$,

(vi) $\displaystyle\sum_{k=0}^{1-a_{i,j}}(-1)^k e_i^{(1-a_{ij}-k)}e_je_i^{(k)} = \sum_{k=0}^{1-a_{i,j}}(-1)^k f_i^{(1-a_{ij}-k)}f_jf_i^{(k)} = 0$ for $i\neq j$.

Here $e_i^{(k)} := e_i^k/[k]_i!$ and $f_i^{(k)} := f_i^k/[k]_i!$ for $k\in\mathbb{Z}_{\geq 0}$.

The quantum group $U_q(\mathfrak{g})$ has a comultiplication Δ which is defined by

$$\Delta(e_i) = e_i\otimes K_i^{-1} + 1\otimes e_i,\quad \Delta(f_i) = f_i\otimes 1 + K_i\otimes f_i \quad\text{and}\quad \Delta(q^h) = q^h\otimes q^h.$$

1.2. **Quiver Hecke algebras.** In this subsection, we shall review symmetric quiver Hecke algebras. Let $\mathbf{k}$ be a field and $\mathsf{A}=(a_{ij})_{i,j\in I}$ be a *symmetric* Cartan matrix. We denote by Q^+ the positive root lattice associated with A. We normalize the symmetric bilinear form $(\ ,\ )$ by $(\alpha_i,\alpha_j)=a_{ij}$. We choose a matrix $\mathcal{Q}^{\mathsf{A}} = (\mathcal{Q}_{ij})_{i,j\in I}$

$$\mathcal{Q}_{i,j}(u,v) = \begin{cases} c_{ij}(u-v)^{-a_{ij}} & \text{if } i\neq j,\\ 0 & \text{if } i=j.\end{cases} \tag{1.1}$$

where $c_{ij}\in\mathbf{k}^\times$ and we assume $\mathcal{Q}_{i,j}(u,v)=\mathcal{Q}_{j,i}(v,u)$.

For $\beta\in\mathsf{Q}^+$ with $\mathrm{ht}(\beta)=n$, we set

$$I^\beta := \{\nu=(\nu_1,\ldots,\nu_n)\in I^n \mid \alpha_{\nu_1}+\ldots+\alpha_{\nu_n}=\beta\}.$$

The symmetric group $\mathfrak{S}_n = \langle s_i \mid 1\leq i<n\rangle$ on n-letters acts on I^n by place permutations in a natural way.

Definition 1.2 ([21, 30])**.** For $\beta \in \mathsf{Q}^+$ with $\mathrm{ht}(\beta) = n$, the *symmetric quiver Hecke algebra* (or KLR-algebra) $R(\beta)$ at β associated with a matrix $\mathcal{Q}^{\mathsf{A}}$ is the $\mathbf{k}$-algebra generated by three sets of generators

$$\{x_k \mid 1 \le k \le n\}, \quad \{\tau_\ell \mid 1 \le \ell < n\}, \quad \{e(\nu) \mid \nu \in I^\beta\}$$

satisfying the following relations:

(i) $e(\nu)e(\nu') = \delta_{\nu,\nu'}e(\nu)$, $\sum_{\nu \in I^\beta} e(\nu) = 1$,

(ii) $x_k e(\nu) = e(\nu) x_k$, $x_k x_{k'} = x_{k'} x_k$,

(iii) $\tau_\ell e(\nu) = e(s_\ell(\nu))\tau_\ell$, $\tau_{\ell'}\tau_\ell = \tau_\ell \tau_{\ell'}$ if $|\ell - \ell'| > 1$,

(iv) $\tau_\ell^2 e(\nu) = \mathcal{Q}_{\nu_\ell,\nu_{\ell+1}}(x_\ell, x_{\ell+1})e(\nu)$,

(v) $$(\tau_\ell x_k - x_{s_\ell(k)}\tau_\ell)e(\nu) = \begin{cases} -e(\nu) & \text{if } k = \ell \text{ and } \nu_\ell = \nu_{\ell+1}, \\ e(\nu) & \text{if } k = \ell+1 \text{ and } \nu_\ell = \nu_{\ell+1}, \\ 0 & \text{otherwise,} \end{cases}$$

(vi) $$(\tau_{\ell+1}\tau_\ell\tau_{\ell+1} - \tau_\ell\tau_{\ell+1}\tau_\ell)e(\nu) = \delta_{\nu_\ell,\nu_{\ell+2}} \frac{\mathcal{Q}_{\nu_\ell,\nu_{\ell+1}}(x_\ell, x_{\ell+1}) - \mathcal{Q}_{\nu_\ell,\nu_{\ell+1}}(x_{\ell+2}, x_{\ell+1})}{x_\ell - x_{\ell+2}} e(\nu).$$

Note that, for each α_i, the quiver Hecke algebra $R(\alpha_i)$ is isomorphic to $\mathbf{k}[x]$. Thus there exists an $R(\alpha_i)$-module $L(i) = \mathbf{k}u(i)$ defined by $x \cdot u(i) = 0$.

We denote by $\mathrm{Rep}(R(\beta))$ the category of finite-dimensional $R(\beta)$-modules M such that the actions of x_k $(1 \le k \le \mathrm{ht}(\beta))$ on M are nilpotent.

For $\beta, \gamma \in \mathsf{Q}^+$ with $\mathrm{ht}(\beta) = m$ and $\mathrm{ht}(\gamma) = n$, we define the idempotent $e(\beta, \gamma)$ in $R(\beta + \gamma)$ by

$$e(\beta, \gamma) = \sum_{\nu \in I^\beta, \mu \in I^\gamma} e(\nu * \mu)$$

where $\nu * \mu = (\nu_1, \ldots, \nu_m, \mu_1, \ldots, \mu_n)$ for $\nu := (\nu_1, \ldots, \nu_m) \in I^\beta$ and $\mu := (\mu_1, \ldots, \mu_n) \in I^\gamma$.

Let $\iota_{\beta,\gamma}$ be a $\mathbf{k}$-algebra homomorphism

$$\iota_{\beta,\gamma} \colon R(\beta) \otimes R(\gamma) \longrightarrow e(\beta, \gamma) R(\beta + \gamma) e(\beta, \gamma)$$

given by

$$\begin{aligned} &\iota_{\beta,\gamma}(e(\nu) \otimes e(\mu)) = e(\nu * \mu), \\ &\iota_{\beta,\gamma}(x_k \otimes 1) = x_k e(\beta, \gamma) \ (1 \le k \le m), \\ &\iota_{\beta,\gamma}(1 \otimes x_k) = x_{m+k} e(\beta, \gamma) \ (1 \le k \le n), \\ &\iota_{\beta,\gamma}(\tau_k \otimes 1) = \tau_k e(\beta, \gamma) \ (1 \le k < m), \\ &\iota_{\beta,\gamma}(1 \otimes \tau_k) = \tau_{m+k} e(\beta, \gamma) \ (1 \le k < n). \end{aligned} \tag{1.2}$$

For an $R(\beta)$-module M and an $R(\gamma)$-module N, we define their *convolution product* $M \circ N$ as follows:

$$M \circ N := R(\beta+\gamma)e(\beta,\gamma) \underset{R(\beta)\otimes R(\gamma)}{\otimes} (M \otimes N). \tag{1.3}$$

Then the category $\mathrm{Rep}(R) := \bigoplus_{\beta \in \mathsf{Q}^+} \mathrm{Rep}(R(\beta))$ becomes a tensor category in the sense of [15, Appendix A.1] induced by the convolution product. We denote by $\mathrm{K}(\mathrm{Rep}(R))$ the Grothendieck ring of the category $\mathrm{Rep}(R)$.

1.3. **Quantum affine algebras.** In this subsection, we briefly recall the representation theory of the quantum affine algebra $U_q'(\mathfrak{g})$. We refer to [1, 4, 20] for the basics on the representation theory of quantum affine algebra $U_q'(\mathfrak{g})$.

The affine Cartan datum $(\mathsf{A}, \mathsf{P}, \mathsf{P}^\vee, \Pi, \Pi^\vee)$ consists of

(1) an affine Cartan matrix A,
(2) a weight lattice $\mathsf{P} = (\bigoplus_{i\in I} \mathbb{Z}\Lambda_i) \oplus \mathbb{Z}\delta$, where δ is the *imaginary root*,
(3) a set of simple roots $\Pi = \{\alpha_i \mid i \in I\}$, where

$$\alpha_i = \begin{cases} \sum_{j\in I} a_{ji}\Lambda_j & \text{if } i \in I_0, \\ \sum_{j\in I} a_{j0}\Lambda_j + \delta & \text{if } i = 0, \end{cases}$$

(4) The simple coroots $h_i \in \mathsf{P}^\vee$ are defined by $\langle h_i, \Lambda_j\rangle = \delta_{i,j}$ and $\langle h_i, \delta\rangle = 0$.

Note that

$$\mathbb{Z}\delta = \{\lambda \in \mathsf{Q} \mid \langle h_i, \lambda\rangle = 0 \text{ for all } i \in I\}.$$

Let us denote by $c = \sum_{i\in I} c_i h_i$ $(c_i \in \mathbb{Z}_{>0})$ the unique element in $\mathsf{P}^\vee$ such that

$$\mathbb{Z}c = \{h \in \mathsf{P}^\vee \mid \langle h, \alpha_i\rangle = 0 \text{ for all } i \in I\}.$$

Write $\delta = \sum_{i\in I} d_i\alpha_i$. We take a symmetric bilinear form on $\mathbb{Q}\otimes_{\mathbb{Z}} \mathsf{P}$ such that

$$(\alpha_i, \alpha_j) = c_i d_i^{-1} a_{ij} \quad (i, j \in I) \quad \text{and } \langle h_i, \lambda\rangle = \frac{2(\alpha_i,\lambda)}{(\alpha_i,\alpha_i)} \text{ for all } \lambda \in \mathsf{P} \text{ and } i \in I.$$

Then we have $(\delta, \lambda) = \langle c, \lambda\rangle$ for all $\lambda \in \mathsf{P}$.

We denote by

- $\mathfrak{g}$ the affine Kac-Moody Lie algebra with an affine Cartan datum $(\mathsf{A}, \mathsf{P}, \mathsf{P}^\vee, \Pi, \Pi^\vee)$,
- $U_q(\mathfrak{g})$ the quantum group associated with the affine Cartan datum $(\mathsf{A}, \mathsf{P}, \mathsf{P}^\vee, \Pi, \Pi^\vee)$,
- $U_q'(\mathfrak{g})$ the subalgebra of $U_q(\mathfrak{g})$ generated by e_i, f_i and $K_i^{\pm 1}$ for all $i \in I$.

We call $U'_q(\mathfrak{g})$ the *quantum affine algebra.*

Whenever we deal with the $U'_q(\mathfrak{g})$-modules, we take the base field $\mathbf{k}$ to be the algebraic closure of $\mathbb{C}(q)$ in $\cup_{m>0}\mathbb{C}((q^{1/m}))$.

We take an element 0 in I where 0 denotes the leftmost vertices in the tables in [14, pages 54, 55] except $A^{(2)}_{2n}$-case in which we take the longest simple root as α_0. We set $I_0 := I \setminus \{0\}$.

Let $\mathfrak{g}_0$ be the finite-dimensional simple Lie subalgebra of $\mathfrak{g}$ generated by e_i, f_i for $i \in I_0$, called the *classical part* of $\mathfrak{g}$,

Set $\mathsf{P}_{\rm cl} := \mathsf{P}/\mathbb{Z}\delta$ and call it the *classical weight lattice.* Considering the canonical projection $\mathrm{cl}\colon \mathsf{P} \to \mathsf{P}_{\rm cl}$, we have

$$\mathsf{P}_{\rm cl} = \bigoplus_{i\in I} \mathbb{Z}\mathrm{cl}(\Lambda_i).$$

We say that a $U'_q(\mathfrak{g})$-module M is *integrable* if

(i) it is $\mathsf{P}_{\rm cl}$-graded; i.e.,

$$M = \bigoplus_{\lambda\in\mathsf{P}_{\rm cl}} M_\lambda \quad \text{where } M_\lambda = \{u \in M \mid K_i u = q_i^{\langle h_i,\lambda\rangle} u \ \text{ for all } i \in I\},$$

(ii) for all $i \in I$, e_i and f_i act on M locally nilpotently.

We denote by $\mathscr{C}_\mathfrak{g}$ the category of finite-dimensional integrable $U'_q(\mathfrak{g})$-modules. Note that $\mathscr{C}_\mathfrak{g}$ is a tensor category induced by the comultiplication

$$\Delta|_{U'_q(\mathfrak{g})}\colon U'_q(\mathfrak{g}) \to U'_q(\mathfrak{g}) \otimes U'_q(\mathfrak{g}).$$

A simple module M in $\mathscr{C}_\mathfrak{g}$ contains a non-zero vector u of weight $\lambda \in \mathsf{P}_{\rm cl}$ such that

- $\langle c, \lambda\rangle = 0$ and $\langle h_i, \lambda\rangle \geq 0$ for all $i \in I_0$,
- all the weight of M are contained in $\lambda - \sum_{i\in I_0} \mathbb{Z}_{\geq 0}\mathrm{cl}(\alpha_i)$.

Such a λ is unique and u is unique up to a constant multiple. We call λ the *dominant extremal weight* of M and u the *dominant extremal weight vector* of M.

For a $\mathsf{P}_{\rm cl}$-graded $U'_q(\mathfrak{g})$-module M, $M_{\rm aff}$ denotes the P-graded $U'_q(\mathfrak{g})$-module defined by

(i) $M_{\rm aff} = \bigoplus_{\lambda\in\mathsf{P}} (M_{\rm aff})_\lambda$ such that $(M_{\rm aff})_\lambda \simeq (M)_{\mathrm{cl}(\lambda)}$,

(ii) the actions of e_i and f_i are defined in such a way that they have weight $\pm\alpha_i$ and commute with the canonical projection $\mathrm{cl}\colon M_{\rm aff} \to M$.

For $x \in \mathbf{k}^\times$ and $M \in \mathscr{C}_\mathfrak{g}$, we define

$$M_x := M_{\rm aff}/(z_M - x)M_{\rm aff} \tag{1.4}$$

where z_M denotes the $U_q'(\mathfrak{g})$-module automorphism of M_{aff} of weight δ, defined as the composition $(M_{\mathrm{aff}})_\lambda \xrightarrow{\sim} M_{\mathrm{cl}(\lambda)} \xrightarrow{\sim} (M_{\mathrm{aff}})_{\lambda+\delta}$. For each $i \in I_0$, we set

$$\varpi_i := \gcd(c_0, c_i)^{-1}\mathrm{cl}(c_0\Lambda_i - c_i\Lambda_0) \in \mathsf{P}_{\mathrm{cl}}.$$

Then there exists a unique simple $U_q'(\mathfrak{g})$-module $V(\varpi_i)$ in $\mathscr{C}_\mathfrak{g}$, called the *fundamental module of weight* ϖ_i, satisfying the following conditions:

(i) all the weights of $V(\varpi_i)$ are contained in the convex hull of $W_0\varpi_i$,

(ii) $\dim V(\varpi_i)_{\varpi_i} = 1$ and $V(\varpi_i) = U_q'(\mathfrak{g}) \cdot u_{\varpi_i}$,

(iii) for any $\mu \in W_0\varpi_i \subset \mathsf{P}_{\mathrm{cl}}$, we can associate a non-zero vector u_μ of weight μ such that

$$u_{s_j(\mu)} = \begin{cases} f_j^{(\langle h_j,\mu\rangle)} u_\mu & \text{if } \langle h_j,\mu\rangle \ge 0 \\ e_j^{(-\langle h_j,\mu\rangle)} u_\mu & \text{if } \langle h_j,\mu\rangle \le 0 \end{cases} \quad \text{for any } j \in I.$$

Here W_0 denotes the Weyl group of $\mathfrak{g}_0$ and u_{ϖ_i} denotes the dominant extremal weight vector of weight ϖ_i.

The module $V(\varpi_i)_x$ ($x \in \mathbf{k}^\times$) has the left dual $\big(V(\varpi_i)_x\big)^*$ and the right dual ${}^*\big(V(\varpi_i)_x\big)$ with the following $U_q'(\mathfrak{g})$-module homomorphisms (see [1, §1.3]):

$$\big(V(\varpi_i)_x\big)^* \otimes V(\varpi_i)_x \xrightarrow{\mathrm{tr}} \mathbf{k} \quad \text{and} \quad V(\varpi_i)_x \otimes {}^*\big(V(\varpi_i)_x\big) \xrightarrow{\mathrm{tr}} \mathbf{k},$$

where

$$\text{(1.5)} \quad \big(V(\varpi_i)_x\big)^* \simeq V(\varpi_{i^*})_{x(p^*)^{-1}}, \ {}^*\big(V(\varpi_i)_x\big) \simeq V(\varpi_{i^*})_{xp^*} \quad \text{and} \quad p^* := (-1)^{\langle \rho^\vee, \delta\rangle} q^{(\rho,\delta)}.$$

Here ρ (respectively, $\rho^\vee$) denotes an element in P (respectively, $\mathsf{P}^\vee$) such that $\langle h_i, \rho\rangle = 1$ (respectively, $\langle \rho^\vee, \alpha_i\rangle = 1$) for all $i \in I$, and $i \mapsto i^*$ denotes the involution on I_0 given by $\alpha_i = -w_0\,\alpha_{i^*}$, where w_0 is the longest element of W_0.

We say that a $U_q'(\mathfrak{g})$-module M is *good* if it has a *bar involution*, a crystal basis with *simple crystal graph*, and a *global basis* (see [20] for precise definition). For instance, every fundamental module $V(\varpi_i)$ for $i \in I_0$ is a good module. Note that every good module is a simple $U_q'(\mathfrak{g})$-module. Moreover the tensor product of good modules is again good. Hence any good module M is *real* simple, i.e., $M \otimes M$ is simple.

2. KLR-type Schur-Weyl duality functors and Auslander-Reiten quiver

In this section, we recall the *KLR-type Schur-Weyl duality functor* $\mathcal{F}$ constructed in [15] which depends on the choice of a family of good $U_q'(\mathfrak{g})$-modules and spectral parameters.

We also recall the notion of Auslander-Reiten quiver Γ_Q which was used in [16] to construct an exact functor $\mathcal{F}_Q^{(1)}$ when $\mathfrak{g} = A_N^{(1)}$ and $D_N^{(1)}$.

2.1. **KLR-type Schur-Weyl duality functors.** For simple $U_q'(\mathfrak{g})$-modules M_1 and M_2, there exists a unique $U_q'(\mathfrak{g})$-module homomorphism

$$R^{\mathrm{norm}}_{M_1,M_2}(z_1,z_2) : (M_1)_{\mathrm{aff}} \otimes (M_2)_{\mathrm{aff}} \to \mathbf{k}(z_1,z_2) \underset{\mathbf{k}[z_1^{\pm1},z_2^{\pm1}]}{\otimes} (M_2)_{\mathrm{aff}} \otimes (M_1)_{\mathrm{aff}}$$

which sends $u_1 \otimes u_s$ to $u_2 \otimes u_1$. Here $z_k := z_{V_k}$ denotes the $U_q'(\mathfrak{g})$-automorphism of $(M_k)_{\mathrm{aff}}$ of weight δ, and u_k is a dominant extremal weight vector of M_k $(k = 1, 2)$. We call the $U_q'(\mathfrak{g})$-homomorphism $R^{\mathrm{norm}}_{M_1,M_2}(z_1,z_2)$ the *normalized R-matrix* between M_1 and M_2.

We denote by $d_{M_1,M_2}(z)$ the *denominator* of $R^{\mathrm{norm}}_{M_1,M_2}(z_1,z_2)$ which is the monic polynomial of the smallest degree in $\mathbf{k}[z]$ such that

$$d_{M_1,M_2}(z_2/z_1) R^{\mathrm{norm}}_{M_1,M_2}(z_1,z_2) \colon (M_1)_{\mathrm{aff}} \otimes (M_2)_{\mathrm{aff}} \to (M_2)_{\mathrm{aff}} \otimes (M_1)_{\mathrm{aff}}. \tag{2.1}$$

Let $\{V_s\}_{s\in\mathcal{S}}$ be a family of good $U_q'(\mathfrak{g})$-modules labeled by an index set $\mathcal{S}$. Let u_s be a dominant extremal weight vector of V_s.

For a triple (J, X, s) consisting of an index set J and two maps $X\colon J \to \mathbf{k}^\times$, $s\colon J \to \mathcal{S}$, we define a quiver $\mathscr{S}^J = (\mathscr{S}^J_0, \mathscr{S}^J_1)$ in the following way:

(2.2)
(i) $\mathscr{S}^J_0 = J$.
(ii) For $i, j \in J$, we put d_{ij} many arrows from i to j, where d_{ij} is the order of the zero of $d_{V_{s(i)},V_{s(j)}}(z)$ at $z = X(j)/X(i)$.

Then $\mathscr{S}^J$ is a quiver without loops and 2-cycles, i.e., $\mathsf{d}_{ij}\mathsf{d}_{ji} = 0$. We define a symmetric Cartan matrix $\mathsf{A}^J = (a_{ij})_{i,j\in J}$ by

$$a_{ij} = -\mathsf{d}_{ij} - \mathsf{d}_{ji} \text{ if } i \neq j, \text{ and } a_{ii} = 2.$$

Let Q^+ be the positive root lattice associated with the Cartan matrix A^J. Then, for $\beta \in \mathsf{Q}^+$, we can define the symmetric quiver Hecke algebra $R^J(\beta)$ associated with a matrix $\mathcal{Q}^{\mathsf{A}^J} = (\mathcal{Q}_{ij})_{i,j\in J}$ such that

$$\mathcal{Q}_{ij}(u,v) = (u-v)^{\mathsf{d}_{ij}}(v-u)^{\mathsf{d}_{ji}} \quad \text{for all } i \neq j \in J.$$

Theorem 2.1 ([15])**.** *For each $\beta \in \mathsf{Q}^+$, there exists a functor*

$$\mathcal{F}_\beta \colon \mathrm{Rep}(R^J(\beta)) \to \mathscr{C}_{\mathfrak{g}}$$

enjoying the following properties:

(a) $\mathcal{F}_{\alpha_i}(L(i)) \simeq (V_{s(i)})_{X(i)}$ *for every* $i \in J$, $\mathcal{F}_0(R^J(0)) \simeq \mathbf{k}$ *for every* $n \in \mathbb{Z}$.

(b) *Set*

$$\mathcal{F} := \bigoplus_{\beta \in \mathsf{Q}^+} \mathcal{F}_\beta \colon \mathrm{Rep}(R^J) \to \mathscr{C}_{\mathfrak{g}}.$$

Then for any $M \in \mathrm{Rep}(R^J(\beta))$ *and* $N \in \mathrm{Rep}(R^J(\gamma))$, *there exists an* $U_q'(\mathfrak{g})$*-module isomorphism*

$$\mathcal{F}(M \circ N) \simeq \mathcal{F}(M) \otimes \mathcal{F}(N)$$

functorial in M *and* N.

(c) *If the underlying graph of a quiver* $\mathscr{S}^J$ *is of finite type* A, D *or* E, *then the functor* $\mathcal{F}$ *is exact. In particular,* $\mathcal{F}$ *induces a ring homomorphism*

$$\mathrm{K}(\mathrm{Rep}(R^J)) \to \mathrm{K}(\mathscr{C}_{\mathfrak{g}}).$$

We call $\mathcal{F}$ the *KLR-type Schur-Weyl duality functor*.

2.2. **Auslander-Reiten quiver Γ_Q.** In this subsection, we briefly review the combinatorial feature of the Auslander-Reiten quiver Γ_Q of a Dynkin quiver Q of a finite-dimensional simply-laced simple Lie algebra $\mathfrak{g}_0$.

Throughout this subsection, we denote by w_0 the longest element of the Weyl group W_0 of $\mathfrak{g}_0$, $\Pi_{\mathfrak{g}_0}$ the set of simple roots of $\mathfrak{g}_0$ and $\Delta^+_{\mathfrak{g}_0}$ (respectively, $\Delta^-_{\mathfrak{g}_0}$) the set of positive (respectively, negative) roots of $\mathfrak{g}_0$.

Let $Q = (Q_0, Q_1)$ be a Dynkin quiver of $\mathfrak{g}_0$. For a vertex i, we denote by s_iQ the quiver obtained from Q by reversing all arrows in Q_1 incident to i. We say that a vertex $i \in Q_0$ is a *source* (respectively, *sink*) if every arrow incident to i starts (respectively, ends) at i. For a reduced expression $\widetilde{w} = s_{i_1} \cdots s_{i_l}$ of $w \in W_0$, we say that $\widetilde{w}$ is *adapted to* Q if

$$i_k \text{ is a source of the quiver } s_{i_{k-1}} \cdots s_{i_1} Q \text{ for all } 1 \le k \le l.$$

It is known that there exists a unique Coxeter element τ which has a reduced expression adapted to Q. In fact, all the reduced expressions of such a τ are adapted to Q.

A map $\xi \colon Q_0 \to \mathbb{Z}$ is called a *height function* if $\xi_i = \xi_j + 1$ if there exists an arrow $i \to j \in Q_1$. Since Q is connected, any pair ξ and ξ' of height functions differ by a constant. We fix such a height function and we define the repetition quiver $\widehat{Q}$ as follows:

(i) $\widehat{Q}_0 = \{(i, p) \in Q_0 \times \mathbb{Z} \mid p - \xi_i \in 2\mathbb{Z}\}$,

(ii) $\widehat{Q}_1 = \{(i, p) \to (j, p+1) \mid i \text{ and } j \text{ are adjacent in the Dynkin diagram of } \mathfrak{g}_0\}$.

Note that there is a path from (i,p) to (j,r) in $\widehat{Q}$ if and only if $d(i,j) \le r-p$, where $d(i,j)$ is the distance of i and j in the Dynkin diagram except the case $\mathfrak{g}_0 \simeq A_1$.

Set $\widehat{\Delta}_{\mathfrak{g}_0} := \Delta^+_{\mathfrak{g}_0} \times \mathbb{Z}$. For $i \in I_0$, we define an element $\gamma_i \in \Delta^+_{\mathfrak{g}_0}$ as follows:

$$\gamma_i = \sum_{j \in B(i)} \alpha_j, \tag{2.3}$$

where $B(i)$ denotes the set of vertices j such that there exists a path from j to i in Q.

We define a bijection $\phi\colon \widehat{Q}_0 \to \widehat{\Delta}_{\mathfrak{g}_0}$ in an inductive way ([12, §2.2]):

(2.4)

(i) $\phi(i,\xi_i) = (\gamma_i, 0)$,

(ii) for a given $\beta \in \Delta^+_{\mathfrak{g}_0}$ with $\phi(i,p) = (\beta, m)$,

- if $\tau(\beta) \in \Delta^+_{\mathfrak{g}_0}$, then $\phi(i,p-2) := (\tau(\beta), m)$,
- if $\tau(\beta) \in \Delta^-_{\mathfrak{g}_0}$, then $\phi(i,p-2) := (-\tau(\beta), m-1)$,
- if $\tau^{-1}(\beta) \in \Delta^+_{\mathfrak{g}_0}$, then $\phi(i,p+2) := (\tau^{-1}(\beta), m)$,
- if $\tau^{-1}(\beta) \in \Delta^-_{\mathfrak{g}_0}$, then $\phi(i,p+2) := (-\tau^{-1}(\beta), m+1)$.

The *Auslander-Reiten quiver* Γ_Q is the full subquiver of $\widehat{Q}$ whose set of vertices is $\phi^{-1}(\Delta^+_{\mathfrak{g}_0} \times \{0\})$.

We say that a (partial) order $\preceq$ on $\Delta^+_{\mathfrak{g}_0}$ is *convex*, if it satisfies the condition:

(2.5) either $\beta \preceq \beta+\gamma \preceq \gamma$ or $\gamma \preceq \beta+\gamma \preceq \beta$ whenever $\beta, \gamma \in \Delta^+_{\mathfrak{g}_0}$ satisfy $\beta+\gamma \in \Delta^+_{\mathfrak{g}_0}$.

The Auslander-Reiten quiver Γ_Q defines a convex order $\preceq_Q$ on $\Delta^+_{\mathfrak{g}_0}$ as follows (cf. [29]):

$$\beta \preceq_Q \gamma \text{ if and only if } d(i,j) \le a-b, \tag{2.6}$$

where $\phi^{-1}(\beta,0) = (i,a)$ and $\phi^{-1}(\gamma,0) = (j,b)$. Note that ([2, 29], see also [33, §1])

$$\beta \preceq_Q \gamma \text{ if and only if there exists a path from } \phi^{-1}(\gamma,0) \text{ to } \phi^{-1}(\beta,0) \text{ in } \Gamma_Q. \tag{2.7}$$

On the other hand, for an arbitrary finite-dimensional simple Lie algebra $\mathfrak{g}_0$, any reduced expression $\widetilde{w_0} = s_{i_1} \cdots s_{i_r}$ of the longest element w_0 of $\mathfrak{g}_0$ induces a convex total order $\le_{\widetilde{w_0}}$ on $\Delta^+_{\mathfrak{g}_0}$

$$\beta_k \le_{\widetilde{w_0}} \beta_l \text{ if and only if } k \le l.$$

Here $\beta_k := s_{i_1} \cdots s_{i_{k-1}} \alpha_{i_k}$ $(1 \le k \le r)$. Conversely, any convex total order is obtained in this way by a reduced expression of w_0 ([28]).

It is proved in [2] that, for every reduced expression $\widetilde{w_0}$ of w_0 adapted to Q, the convex total order induced by $\widetilde{w_0}$ is stronger than the order $\preceq_Q$, i.e., for $\beta, \gamma \in \Delta^+_{\mathfrak{g}_0}$, $\beta \preceq_Q \gamma$ implies $\beta \leq_{\widetilde{w_0}} \gamma$.

For a convex order $\preceq$ and a positive root α, a pair of positive roots (β, γ) is called a *minimal pair of* α if $\beta + \gamma = \alpha$, $\beta \preceq \alpha \preceq \gamma$ and there exists no pair (β', γ') other than (β, γ) such that $\beta' + \gamma' = \alpha$ and $\beta \preceq \beta' \preceq \alpha \preceq \gamma' \preceq \gamma$.

2.3. The functor $\mathcal{F}^{(1)}_Q$.

In this subsection, we briefly review the properties of the functor $\mathcal{F}^{(1)}_Q$ which were investigated in [16].

For simplicity, we denote by $d_{k,l}(z)$ or $d^{\mathfrak{g}}_{k,l}(z)$ the denominator $d_{V(\varpi_k),V(\varpi_l)}(z)$ between the fundamental $U'_q(\mathfrak{g})$-modules $V(\varpi_k)$ and $V(\varpi_l)$ $(k, l \in I_0)$ (see (2.1)).

In [1, 16], the denominators $d_{k,l}(z)$ for $\mathfrak{g} = A^{(1)}_N$ and $D^{(1)}_N$ are computed as follows. Note that we have $d_{k,l}(z) = d_{l,k}(z)$.

Theorem 2.2.

(a) *For* $\mathfrak{g} = A^{(1)}_N$ $(N \geq 2)$ *and* $1 \leq k, l \leq N$, *we have*

$$d^{A^{(1)}_N}_{k,l}(z) = \prod_{s=1}^{\min(k,l,N+1-k,N+1-l)} (z - (-q)^{|k-l|+2s}). \tag{2.8}$$

(b) *For* $\mathfrak{g} = D^{(1)}_N$ $(N \geq 4)$ *and* $1 \leq k, l \leq N$, *we have*

$$d^{D^{(1)}_N}_{k,l}(z) = \begin{cases} \displaystyle\prod_{s=1}^{\min(k,l)} (z - (-q)^{|k-l|+2s})(z - (-q)^{2N-2-k-l+2s}) & \text{if } 1 \leq k, l \leq N-2, \\ \displaystyle\prod_{s=1}^{k} (z - (-q)^{N-k-1+2s}) & \text{if } 1 \leq k \leq N-2 \text{ and } l \in \{N-1, N\}, \\ \displaystyle\prod_{s=1}^{\lfloor \frac{N-1}{2} \rfloor} (z - (-q)^{4s}) & \text{if } k \neq l \in \{N, N-1\}, \\ \displaystyle\prod_{s=1}^{\lfloor \frac{N}{2} \rfloor} (z - (-q)^{4s-2}) & \text{if } k = l \in \{N-1, N\}. \end{cases} \tag{2.9}$$

The following theorem is one of the main results in [16].

Theorem 2.3 ([16, Theorem 4.3.1]). *Let $\mathfrak{g}^{(1)}$ be an affine Kac-Moody algebra of type $A_N^{(1)}$ or $D_N^{(1)}$. We take the index set*

$$J := \phi^{-1}(\Pi_{\mathfrak{g}_0^{(1)}} \times \{0\})$$

and two maps $s^{(1)} \colon J \to I_0^{(1)} := \{1, 2, \dots, N\}$ and $X^{(1)} \colon J \to \mathbf{k}^\times$ such that

$$s^{(1)}(i,p) = i \quad \text{and} \quad X^{(1)}(i,p) = (-q)^p.$$

Then the quiver $\mathscr{S}^J$ in (2.2) coincides with the quiver Q^{rev} obtained from Q by reversing all the arrows.

For a given Dynkin quiver Q of $\mathfrak{g}_0^{(1)}$, we set

$$V_Q(\beta) := V(\varpi_i)_{(-q)^p} \quad \text{where } \phi^{-1}(\beta, 0) = (i,p).$$

Definition 2.4 ([12]). The category $\mathscr{C}_Q^{(1)}$ is defined to be the smallest full subcategory of $\mathscr{C}_{\mathfrak{g}^{(1)}}$ such that

(a) it contains $V_Q(\beta)$ for all $\beta \in \Delta_{\mathfrak{g}_0}^+$ and the trivial representation,
(b) it is stable under taking tensor products, subquotients and extensions.

We define an order $\prec_{\mathbb{Z}}$ on $\mathbb{Z}_{\geq 0}^r$ ($r \in \mathbb{Z}_{>0}$) as follows:

(2.10) $\underline{m}' = (m'_1, \dots, m'_r) \prec_{\mathbb{Z}} \underline{m} = (m_1, \dots, m_r)$ if and only if there exist integers k, s such that $1 \leq k \leq s \leq r$, $m'_t = m_t$ $(t < k)$, $m'_k < m_k$, and $m'_t = m_t$ $(s < t \leq r)$, $m'_s < m_s$.

Theorem 2.5 ([24], see also [22]). *Let $\mathfrak{g}_0$ be a finite-dimensional simple Lie algebra. We fix a reduced expression $\widetilde{w_0} = s_{i_1} \cdots s_{i_r}$ of the longest element $w_0 \in W_0$, and set $\beta_k = s_{i_1} \cdots s_{i_{k-1}} \alpha_{i_k}$. Let R be the quiver Hecke algebra corresponding to $\mathfrak{g}_0$. Then for every $\beta_k \in \Delta_{\mathfrak{g}_0}^+$, there exists a simple $R(\beta_k)$-module $S_{\widetilde{w_0}}(\beta_k)$ satisfying the following properties:*

(i) *For every $\underline{m} = (m_1, \dots, m_r) \in \mathbb{Z}_{\geq 0}^r$, there exists a non-zero R-module homomorphism*

$$\mathbf{r}_{\underline{m}} \colon \overrightarrow{\nabla}_{\widetilde{w_0}}(\underline{m}) := S_{\widetilde{w_0}}(\beta_1)^{\circ m_1} \circ \cdots \circ S_{\widetilde{w_0}}(\beta_r)^{\circ m_r} \\ \longrightarrow \overleftarrow{\nabla}_{\widetilde{w_0}}(\underline{m}) := S_{\widetilde{w_0}}(\beta_r)^{\circ m_r} \circ \cdots \circ S_{\widetilde{w_0}}(\beta_1)^{\circ m_1}.$$

Such a morphism is unique up to a constant multiple, and $\mathrm{Im}(\mathbf{r}_{\underline{m}})$ is a simple module and coincides with the head of $\overrightarrow{\nabla}_{\widetilde{w_0}}(\underline{m})$ and the socle of $\overleftarrow{\nabla}_{\widetilde{w_0}}(\underline{m})$.

(ii) *For every* $\underline{m} = (m_1, \dots, m_r) \in \mathbb{Z}^r_{\ge 0}$, *we have*

$$[\overrightarrow{\nabla}_{\widetilde{w_0}}(\underline{m})] \in [\mathrm{Im}(\mathbf{r}_{\underline{m}})] + \sum_{\underline{m}' \prec_{\mathbb{Z}} \underline{m}} \mathbb{Z}_{\ge 0}[\mathrm{Im}(\mathbf{r}_{\underline{m}'})].$$

(iii) *For any simple* $R(\beta)$*-module* M, *there exists a unique* $\underline{m} = (m_1, \dots, m_r) \in \mathbb{Z}^r_{\ge 0}$ *such that*

$$M \simeq \mathrm{Im}(\mathbf{r}_{\underline{m}}) \simeq \mathrm{hd}\big(\overrightarrow{\nabla}_{\widetilde{w_0}}(\underline{m})\big).$$

(iv) *For any minimal pair* (β_k, β_l) *of* $\beta_j = \beta_k + \beta_l$, *there is an exact sequence*

$$0 \to S_{\widetilde{w_0}}(\beta_j) \to S_{\widetilde{w_0}}(\beta_k) \circ S_{\widetilde{w_0}}(\beta_l) \xrightarrow{\mathbf{r}} S_{\widetilde{w_0}}(\beta_l) \circ S_{\widetilde{w_0}}(\beta_k) \to S_{\widetilde{w_0}}(\beta_j) \to 0$$

with a non-zero $\mathbf{r}$.

Remark 2.6. In [24] (see also [22]), it is shown that the simple module $S_{\widetilde{w_0}}(\beta)$ corresponds to the *dual PBW generator* of $U_q^-(\mathfrak{g}_0)$ of weight $-\beta$, under the isomorphism between the Grothendieck ring K(R-gmod) and the dual integral form of $U_q^-(\mathfrak{g}_0)$. Here R-gmod is the category of finite-dimensional *graded* R-modules. As a result, we have the following properties:

(a) For every simple root α_i, we have $S_{\widetilde{w_0}}(\alpha_i) \simeq L(i)$.

(b) For any pair of reduced expressions $\widetilde{w_0}$ and $\widetilde{w_0}'$ of w_0 which are adapted to Q, we have

$$S_{\widetilde{w_0}}(\beta) \simeq S_{\widetilde{w_0}'}(\beta) \quad \text{for all } \beta \in \Delta^+_{\mathfrak{g}_0}.$$

Thus we will denote by $S_Q(\beta)$ the simple $R(\beta)$-module $S_{\widetilde{w_0}}(\beta)$ for a reduced expression $\widetilde{w_0}$ adapted to Q.

From now on, we assume that $\mathfrak{g}^{(1)} = A_N^{(1)}$ or $D_N^{(1)}$, and take a Dynkin quiver Q of type $\mathfrak{g}_0^{(1)}$. We also denote by $R(\beta)$ the quiver Hecke algebra $R^J(\beta)$ associated with the triple $(J, X^{(1)}, s^{(1)})$ in Theorem 2.3. Hence the quiver Hecke algebra is of type A_N or D_N. Then we have an exact functor

$$\mathcal{F}_Q^{(1)} : \mathrm{Rep}(R) \to \mathscr{C}_{\mathfrak{g}^{(1)}}.$$

By Theorem 2.1 (a), one can observe that

$$\mathcal{F}_Q^{(1)}(L(j)) \simeq V_Q(\alpha_j) \quad \text{for any } j \in I_0.$$

Theorem 2.7 ([16]).

(i) *For every* $\beta \in \Delta^+_{\mathfrak{g}_0}$, *we have*

$$\mathcal{F}_Q^{(1)}(S_Q(\beta)) \simeq V_Q(\beta).$$

(ii) *The functor* $\mathcal{F}_Q^{(1)}$ *sends a simple object to a simple object and it induces a ring isomorphism*

$$\varphi^{(1)} \colon \mathrm{K}(\mathrm{Rep}(R)) \xrightarrow{\sim} \mathrm{K}(\mathscr{C}_Q^{(1)}).$$

3. Similarities between untwisted cases and twisted cases

In this section, we first recall the denominators of normalized R-matrices of type $A_N^{(2)}$ and $D_N^{(2)}$. Next, we shall observe similarities between the representations of quantum affine algebras of untwisted types and twisted types, i.e., the similarity between $A_N^{(1)}$ and $A_N^{(2)}$ $(N \geq 2)$ and the similarity between $D_N^{(1)}$ and $D_N^{(2)}$ $(N \geq 4)$.

Remark 3.1. In this paper, we follow the enumeration of vertices of Dynkin diagram of type $A_N^{(t)}$ $(t = 1, 2,\ N \geq 2)$ and $D_N^{(t)}$ $(t = 1, 2,\ N \geq 4)$ as follows.

- $A_N^{(1)}$ $(N \geq 2)$



- $A_{2n}^{(2)}$ $(n \geq 2)$



$q = q_i$ for $1 \leq i < n$.

- $A_2^{(2)}$



$q_0 = q^2$, $q_1 = q^{1/2}$.

- $A_{2n-1}^{(2)}$ $(n \geq 3)$



$q = q_i$ for $i \neq n$.

- $A_3^{(2)}$



$q = q_0 = q_1$, $q_2 = q^2$.

- $D_N^{(1)}$ $(N \geq 4)$



- $D_N^{(2)}$ $(N \geq 4)$



$q = q_0 = q_{N-1}$.

In particular, the affine Kac-Moody algebra of type $A_3^{(2)}$ should be understood as $D_3^{(2)}$. In these cases, i^* and p^* in (1.5) are given as follows:

$$i^* = \begin{cases} N+1-i & \text{if } \mathfrak{g} = A_N^{(1)}, \\ N-(1-\varepsilon) & \text{if } \mathfrak{g} = D_N^{(1)},\ N \text{ is odd and } i = N-\varepsilon,\ \varepsilon = 0,1, \\ i & \text{otherwise,} \end{cases} \tag{3.1}$$

and

$$p^* = \begin{cases} (-q)^{N+1} & \text{if } \mathfrak{g} = A_N^{(1)}, \\ (-q)^{2N-2} & \text{if } \mathfrak{g} = D_N^{(1)}, \\ -q^{N+1} & \text{if } \mathfrak{g} = A_N^{(2)}, \\ (-1)^N q^{2(N-1)} & \text{if } \mathfrak{g} = D_N^{(2)}. \end{cases} \tag{3.2}$$

In [25], the denominators of normalized R-matrices between fundamental modules of type $A_N^{(2)}$ and $D_N^{(2)}$ are computed as in the following theorem. Note that we have $d_{k,l}(z) = d_{l,k}(z)$.

Theorem 3.2.

(a) *For* $\mathfrak{g} = A_N^{(2)}$ $(N \geq 2)$ *and* $1 \leq k, l \leq \lfloor \frac{N+1}{2} \rfloor$, *we have*

$$d_{k,l}^{A_N^{(2)}}(z) = \prod_{s=1}^{\min(k,l)} (z - (-q)^{|k-l|+2s})(z + q^{N+1}(-q)^{-k-l+2s}). \tag{3.3}$$

(b) *For* $\mathfrak{g} = D_N^{(2)}$ $(N \geq 4)$ *and* $1 \leq k, l \leq N-1$, *we have*

$$d_{k,l}^{D_N^{(2)}}(z) = \begin{cases} \displaystyle\prod_{s=1}^{\min(k,l)} \big(z^2 - (-q^2)^{|k-l|+2s}\big)\big(z^2 - (-q^2)^{2N-2-k-l+2s}\big) & \textit{if } 1 \leq k, l \leq N-2, \\ \displaystyle\prod_{s=1}^{k} \big(z^2 + (-q^2)^{N-1-k+2s}\big) & \textit{if } 1 \leq k < N-1 \textit{ and } l = N-1, \\ \displaystyle\prod_{s=1}^{N-1} \big(z + (-q^2)^{s}\big) & \textit{if } k = l = N-1. \end{cases} \tag{3.4}$$

3.1. **The Quiver $\mathscr{S}_0(\mathfrak{g})$.** For each $\mathfrak{g}$, we define a quiver $\mathscr{S}(\mathfrak{g})$ as follows:

(i) The set of vertices is the set of all equivalence classes $\widehat{I}_{\mathfrak{g}} := (I_0 \times \mathbf{k}^\times)/\sim$, where the equivalence relation is given in the following way:

$$(i, x) \sim (j, y) \quad \text{if and only if} \quad V(\varpi_i)_x \simeq V(\varpi_j)_y. \tag{3.5}$$

(ii) We put d many arrows from (i, x) to (j, y), where d is the order of the zero of $d_{i,j}(z)$ at $z = y/x$.

By [1, (1.7)], for $x, y \in \mathbf{k}^\times$ and $i \in I_0$, and $\mathfrak{g} = A_N^{(t)}$ ($N \geq 2$, $t = 1, 2$) or $\mathfrak{g} = D_N^{(t)}$ ($N \geq 4$, $t = 1, 2$) we have

$$(3.6) \qquad V(\varpi_i)_x \simeq V(\varpi_i)_y \Longleftrightarrow \begin{cases} x^2 = y^2 & \text{if } \mathfrak{g} = A_{2n-1}^{(2)} \ (n \geq 2),\ i = n \\ x^2 = y^2 & \text{if } \mathfrak{g} = D_N^{(2)} \ (N \geq 3),\ 1 \leq i \leq N-2, \\ x = y & \text{otherwise.} \end{cases}$$

Let $\mathscr{S}_0(\mathfrak{g})$ be a connected component of $\mathscr{S}(\mathfrak{g})$. Then $\mathscr{S}_0(\mathfrak{g})$ is uniquely determined up to a spectral parameter and hence $\mathscr{S}_0(\mathfrak{g})$ is unique up to a quiver isomorphism. For the rest of this paper, we take $\mathscr{S}_0(\mathfrak{g})$ as follows:

$$(3.7) \qquad \begin{aligned} &\mathscr{S}_0(A_n^{(1)}) := \{(i, (-q)^p) \in \{1, \ldots, n\} \times \mathbf{k}^\times \mid p \equiv d(1, i) \bmod 2\}, \\ &\mathscr{S}_0(D_{n+1}^{(1)}) := \{(i, (-q)^p) \in \{1, \ldots, n, n+1\} \times \mathbf{k}^\times \mid p \equiv d(n, i) \bmod 2\}, \\ &\mathscr{S}_0(A_{2n}^{(2)}) := \{(i, (-q)^p) \in \{1, \ldots, n\} \times \mathbf{k}^\times \mid p \in \mathbb{Z}\}, \\ &\mathscr{S}_0(A_{2n-1}^{(2)}) := \{(i, \pm(-q)^p) \in \{1, \ldots, n\} \times \mathbf{k}^\times \mid 1 \leq i \leq n,\ p \equiv i+1 \bmod 2\}, \\ &\mathscr{S}_0(D_{n+1}^{(2)}) := \{(i, (\sqrt{-1})q^p), (i', q^{p'}), (n, \pm q^r) \in \{1, \ldots, n\} \times \mathbf{k}^\times \mid 1 \leq i, i' \leq n-1, \\ &\qquad\qquad p \equiv n - i \equiv 0 \bmod 2,\ p' \equiv n - i' \equiv 1 \bmod 2,\ r \equiv 0 \bmod 2\}. \end{aligned}$$

Recall that $d(i, j)$ denotes the distance of i and j in the Dynkin diagram of $\mathfrak{g}_0$.

We define the maps

$$\pi_{A_N^{(1)}} \colon \mathscr{S}(A_N^{(1)}) \to \mathscr{S}(A_N^{(2)}) \quad \text{and} \quad \pi_{D_N^{(1)}} \colon \mathscr{S}(D_N^{(1)}) \to \mathscr{S}(D_N^{(2)})$$

by

$$(3.8) \qquad \pi_{A_N^{(1)}}(i, x) := \begin{cases} (i, x) & \text{if } 1 \leq i \leq \lfloor \dfrac{N+1}{2} \rfloor, \\ (N+1-i, (-1)^N x) & \text{if } \lfloor \dfrac{N+1}{2} \rfloor < i \leq N \end{cases}$$

and

$$(3.9) \quad \pi_{D_N^{(1)}}(i, x) := \begin{cases} \big(i,\ (\sqrt{-1})^{\delta(N \equiv i \bmod 2)+1}\, x\big) \sim \big(i, (\sqrt{-1})^{N-i} x\big) & \text{if } 1 \leq i \leq N-2, \\ \big(N-1, (-1)^i x\big) & \text{if } i \in \{N-1, N\}, \end{cases}$$

for $i \in I_0$ and $x \in \mathbf{k}^\times$.

Proposition 3.3.

(i) *The maps* $\pi_{A_N^{(1)}}$ *and* $\pi_{D_N^{(1)}}$ *are* $2:1$ *quiver morphisms.*

(ii) *They induce quiver isomorphisms*

$$\mathscr{S}_0(A_N^{(1)}) \xrightarrow{\sim} \mathscr{S}_0(A_N^{(2)}) \quad \text{and} \quad \mathscr{S}_0(D_N^{(1)}) \xrightarrow{\sim} \mathscr{S}_0(D_N^{(2)}).$$

(iii) *The maps $\pi_{A_N^{(1)}}$ and $\pi_{D_N^{(1)}}$ commutes with the duality $*$. More precisely, setting $\mathfrak{g}^{(t)} = A_N^{(t)}$ or $D_N^{(t)}$, and denoting by $V^{(t)}(i,x)$ the $U_q'(\mathfrak{g}^{(t)})$-module $V(\varpi_i)_x$, $\big(V^{(1)}(i,x)\big)^* \simeq V^{(1)}(j,y)$ implies $\big(V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(i,x))\big)^* \simeq V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(j,y))$.*

Proof. Our assertions follow from Theorem 2.2, Theorem 3.2, isomorphisms (3.6) and (3.1), (3.2). ◻

By Proposition 3.3, we have the following corollary:

Corollary 3.4. *Let $\mathfrak{g}^{(t)}$ be the affine Kac-Moody algebra of type $A_N^{(t)}$ $(N \geq 2)$ or $D_N^{(2)}$ $(N \geq 4)$ and $\mathfrak{g}_0^{(t)}$ its classical part $(t = 1, 2)$. For any quiver Q of type $\mathfrak{g}_0^{(1)}$ take the index set*

$$J := \phi^{-1}(\Pi_{\mathfrak{g}_0^{(1)}} \times \{0\})$$

and define the two maps $s^{(2)} \colon J \to I_0^{(2)}$, and $X^{(2)} \colon J \to \mathbf{k}^\times$ by

$$\big(s^{(2)}(i,p), X^{(2)}(i,p)\big) = \pi_{\mathfrak{g}^{(1)}}(i, (-q)^p),$$

where $I_0^{(2)}$ denotes the index set of simple roots for $\mathfrak{g}_0^{(2)}$. Then, the quiver $\mathscr{S}^J$ coincides with the quiver Q^{rev} obtained from Q by reversing all the arrows.

By Corollary 3.4, we obtain an exact KLR-type duality functor

$$\mathcal{F}_Q^{(2)} \colon \mathrm{Rep}(R) \to \mathscr{C}_{\mathfrak{g}^{(2)}} \quad \text{for } \mathfrak{g}^{(2)} = A_N^{(2)} \text{ or } D_N^{(2)}. \tag{3.10}$$

Here R denotes the quiver Hecke algebras of type $\mathfrak{g}_0^{(1)} = A_N$ or D_N.

3.2. **Similarities.**

Now we shall observe interesting similarities between the representations of quantum affine algebras of untwisted types and twisted types thorough the maps $\pi_{A_N^{(1)}}$ and $\pi_{D_N^{(1)}}$.

In the sequel, $\mathfrak{g}^{(t)}$ denotes $A_N^{(t)}$ $(N \geq 2)$ or $D_N^{(t)}$ $(N \geq 4)$ for $t = 1, 2$. We denote by $I^{(t)}$ and $I_0^{(t)}$ the index set of simple roots for $\mathfrak{g}^{(t)}$ and its classical part $\mathfrak{g}_0^{(t)}$, respectively. We denote by $V^{(t)}(\varpi_i)$ the fundamental module for $U_q'(\mathfrak{g}^{(t)})$. For $(i,x) \in I_0^{(t)} \times \mathbf{k}^\times$, we set $V^{(t)}(i,x) := V^{(t)}(\varpi_i)_x$.

For any Dynkin quiver Q of type $\mathfrak{g}_0^{(1)}$ and a height functions ξ, we set

$$V_Q^{(1)}(\beta) := V^{(1)}(i,x) \text{ and } V_Q^{(2)}(\beta) := V^{(2)}\big(\pi_{\mathfrak{g}^{(1)}}(i,x)\big)$$

where $\phi^{-1}(\beta, 0) = (i, p)$ and $x = (-q)^p$. Here ϕ is the map in (2.4), and β is a positive root in $\Delta^+_{\mathfrak{g}^{(1)}}$.

Proposition 3.5 ([10])**.** *Let* $\mathfrak{g}^{(1)} = A^{(1)}_N$ *or* $D^{(1)}_N$*. For any* $(i, x) \in \mathscr{S}(\mathfrak{g}^{(1)})$*, we have*

$$\dim_{\mathbf{k}} V^{(1)}(i, x) = \dim_{\mathbf{k}} V^{(2)}(\pi_{\mathfrak{g}}(i, x)).$$

Proposition 3.6 ([5, Theorem 6.1, Theorem 7.1])**.**

(a) *Let* $\mathfrak{g}^{(1)} = A^{(1)}_N$ *and* $(i, x), (j, y), (k, z) \in I^{(1)}_0 \times \mathbf{k}^\times$*. Then*

$$\operatorname{Hom}_{U'_q(\mathfrak{g}^{(1)})}\big(V^{(1)}(i, x) \otimes V^{(1)}(j, y), V^{(1)}(k, z)\big) \neq 0$$

if and only if one of the following conditions holds:

$$\begin{cases} \text{(A-i) } i + j < N + 1,\ k = i + j,\ x/z = (-q)^{-j},\ y/z = (-q)^{i}, \\ \text{(A-ii) } i + j > N + 1,\ k = i + j - N - 1,\ x/z = (-q)^{-N-1+j},\ y/z = (-q)^{N+1-i}. \end{cases} \tag{3.11}$$

(b) *Let* $\mathfrak{g}^{(1)} = D^{(1)}_N$ *and* $(i, x), (j, y), (k, z) \in I^{(1)}_0 \times \mathbf{k}^\times$*. Then*

$$\operatorname{Hom}_{U'_q(\mathfrak{g}^{(1)})}\big(V^{(1)}(i, x) \otimes V^{(1)}(j, y), V^{(1)}(k, z)\big) \neq 0$$

if and only if one of the following conditions holds:

$$\begin{cases} \text{(D-i) } \ell := \max(i, j, k) \le N - 2,\ s + m = \ell \text{ for } \{s, m\} := \{i, j, k\} \setminus \{\ell\} \text{ and} \\ \qquad \big(x/z, y/z\big) = \begin{cases} \big((-q)^{-j}, (-q)^{i}\big), & \text{if } \ell = k, \\ \big((-q)^{-j}, (-q)^{-i+2N-2}\big), & \text{if } \ell = i, \\ \big((-q)^{j-2N+2}, (-q)^{i}\big), & \text{if } \ell = j, \end{cases} \\ \text{(D-ii) } i + j \ge N,\ k = 2N - 2 - i - j,\ \max(i, j, k) \le N - 2, \text{ and } x/z = (-q)^{-j}, \\ \qquad y/z = (-q)^{i}, \\ \text{(D-iii) } s := \min(i, j, k) \le N - 2,\ \{m, \ell\} := \{i, j, k\} \setminus \{s\} \subset \{N - 1, N\} \text{ and} \\ \qquad \bullet\ N - s \equiv \begin{cases} \ell - m \bmod 2 & \text{if } s = k, \\ \ell - m^* \bmod 2 & \text{if } s \neq k, \end{cases} \\ \qquad \bullet\ \big(x/z, y/z\big) = \begin{cases} \big((-q)^{-N+k+1}, (-q)^{N-k-1}\big), & \text{if } s = k, \\ \big((-q)^{-N+i+1}, (-q)^{2i}\big), & \text{if } s = i, \\ \big((-q)^{-2j}, (-q)^{N-j-1}\big), & \text{if } s = j. \end{cases} \end{cases} \tag{3.12}$$

Remark 3.7.

(a) Note that the triples $((i,x),(j,y),(k,z))$ in Proposition 3.6 belong to the same connected component of $\mathscr{S}(\mathfrak{g}^{(1)})$. $\mathfrak{g}^{(1)} = A_N^{(1)}$ or $D_N^{(1)}$.

(b) Let $((i,x),(j,y),(k,z))$ be a triple appearing in Proposition 3.6. Then the normalized R-matrix $R^{\mathrm{norm}}_{V^{(1)}(\varpi_i),V^{(1)}(\varpi_j)}(z)$ has a multiple pole at $z = y/x$ if and only if $((i,x),(j,y),(k,z))$ satisfies the condition (D-ii) in (3.12). (See Lemma 3.2.4 in [16].)

Proposition 3.8 ([25, Section 3], [18, Corollary 3.3.2])**.**

(a) *Let* $(i,x),(j,y),(k,z) \in \mathscr{S}_0(A_N^{(2)})$. *Assume that*

$$(i,x) = \pi_{A_N^{(1)}}(i',x'),\ (j,y) = \pi_{A_N^{(1)}}(j',y') \ \text{and}\ (k,z) = \pi_{A_N^{(1)}}(k',z')$$

such that $(i',x'),(j',y'),(k',z') \in \mathscr{S}_0(A_N^{(1)})$ *satisfy one of the conditions in* (3.11). *Then we have an epimorphism*

$$V^{(2)}(i,x) \otimes V^{(2)}(j,y) \twoheadrightarrow V^{(2)}(k,z).$$

(b) *Let* $(i,x),(j,y),(k,z) \in \mathscr{S}_0(D_N^{(2)})$. *Assume that*

$$(i,x) = \pi_{D_N^{(1)}}(i',x'),\ (j,y) = \pi_{D_N^{(1)}}(j',y') \ \text{and}\ (k,z) = \pi_{D_N^{(1)}}(k',z')$$

such that $(i',x'),(j',y'),(k',z') \in \mathscr{S}_0(D_N^{(1)})$ *satisfy one of the conditions* (D-i) *or* (D-iii) *in* (3.12) *Then we have an epimorphism*

$$V^{(2)}(i,x) \otimes V^{(2)}(j,y) \twoheadrightarrow V^{(2)}(k,z).$$

Proposition 3.9 ([26, 27])**.** *Let* $\mathfrak{g}^{(1)} = A_N^{(1)}$ *(respectively,* $D_N^{(1)}$*), and let* Q *be a Dynkin quiver of type* $\mathfrak{g}_0^{(1)}$. *Assume that* (β,γ) *is a minimal pair of* $\alpha \in \Delta^+_{\mathfrak{g}_0^{(1)}}$ *with respect to the convex total order induced by a reduced expression* $\widetilde{w_0}$ *of* w_0 *adapted to* Q*, and write*

$$\phi^{-1}(\beta,0) = (i,a),\ \phi^{-1}(\gamma,0) = (j,b) \ \text{and}\ \phi^{-1}(\alpha,0) = (k,c).$$

Then the triple $(i,(-q)^a),(j,(-q)^b),(k,(-q)^c)$ *satisfies one of the conditions* (A-i) *or* (A-ii) *in* (3.11) *(respectively,* (D-i) *or* (D-iii) *in* (3.12)*). In particular, there is a surjective homomorphism*

$$V_Q^{(t)}(\gamma) \otimes V_Q^{(t)}(\beta) \twoheadrightarrow V_Q^{(t)}(\alpha).$$

In Section 5, we will prove the following theorem by using the exact functor $\mathcal{F}_Q^{(2)}$. Note that Proposition 3.8 is a special case of this theorem.

Theorem 3.10. *Let* $\mathfrak{g}^{(1)} = A_N^{(1)}$ *or* $D_N^{(1)}$, *and let* $(i,x),(j,y),(k,z) \in \mathscr{S}(\mathfrak{g}^{(1)})$. *Then there exists an epimorphism*

$$V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(i,x)) \otimes V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(j,y)) \twoheadrightarrow V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(k,z))$$

if and only if there exists an epimorphism

$$V^{(1)}(i,x) \otimes V^{(1)}(j,y) \twoheadrightarrow V^{(1)}(k,z).$$

Note that, for an arbitrary quantum affine algebra $U_q'(\mathfrak{g})$, $V(\varpi_i)_x \otimes V(\varpi_j)_y$ has a simple head by [17]. Hence we have

$$\dim \mathrm{Hom}_{U_q'(\mathfrak{g})}\big(V(\varpi_i)_x \otimes V(\varpi_j)_y, V(\varpi_k)_z\big) \leq 1.$$

4. THE FUNCTOR $\mathcal{F}_Q^{(2)}$

In this section, we investigate the properties of the exact functor $\mathcal{F}_Q^{(2)}$ given in (3.10). Recall that R is the symmetric quiver Hecke algebra of type $\mathfrak{g}_0^{(1)}$.

Theorem 4.1. *For any Dynkin quiver Q of type $\mathfrak{g}_0^{(1)}$ and $\alpha \in \Delta^+_{\mathfrak{g}_0^{(1)}}$, we have*

$$\mathcal{F}_Q^{(2)}(S_Q(\alpha)) \simeq V_Q^{(2)}(\alpha),$$

where $S_Q(\alpha)$ denotes the $R(\alpha)$-module defined in Remark 2.6.

Proof. We shall prove our assertion by induction on $\mathrm{ht}(\alpha)$. For α with $\mathrm{ht}(\alpha) = 1$, our assertion follows from Theorem 2.1 (a). Now we assume that $\mathrm{ht}(\alpha) \geq 2$. Note that there exists a minimal pair (β, γ) of α. By Theorem 2.5 (c), we have a six-term exact sequence of $R(\alpha)$-modules

$$0 \to S_Q(\alpha) \longrightarrow S_Q(\beta) \circ S_Q(\gamma) \xrightarrow{r} S_Q(\gamma) \circ S_Q(\beta) \xrightarrow{s} S_Q(\alpha) \to 0.$$

Applying the functor $\mathcal{F}_Q^{(2)}$, we have an exact sequence of $U_q'(\mathfrak{g})$-modules by the induction hypothesis

$$0 \to \mathcal{F}_Q^{(2)}(S_Q(\alpha)) \to V_Q^{(2)}(\beta) \otimes V_Q^{(2)}(\gamma) \xrightarrow{\mathcal{F}_Q^{(2)}(r)} V_Q^{(2)}(\gamma) \otimes V_Q^{(2)}(\beta) \xrightarrow{\mathcal{F}_Q^{(2)}(s)} \mathcal{F}_Q^{(2)}(S_Q(\alpha)) \to 0.$$

On the other hand, Proposition 3.9 implies that $V_Q^{(2)}(\gamma) \otimes V_Q^{(2)}(\beta)$ is not simple.

We have then $\mathcal{F}_Q^{(2)}(S_Q(\alpha)) \neq 0$. Indeed, if it vanished, we would have

$$V_Q^{(2)}(\beta) \otimes V_Q^{(2)}(\gamma) \simeq V_Q^{(2)}(\gamma) \otimes V_Q^{(2)}(\beta),$$

which implies that $V_Q^{(2)}(\gamma) \otimes V_Q^{(2)}(\beta)$ is simple by [17, Corollary 3.16].

Hence $\mathcal{F}_Q^{(2)}(S_Q(\alpha))$ is the image of a non-zero homomorphism $V_Q^{(2)}(\gamma) \otimes V_Q^{(2)}(\beta) \to V_Q^{(2)}(\beta) \otimes V_Q^{(2)}(\gamma)$. Hence, [17] and the quantum affine version of [19, Proposition 4.5] imply that $\mathcal{F}_Q^{(2)}(S_Q(\alpha))$ is the simple head of $V_Q^{(2)}(\gamma) \otimes V_Q^{(2)}(\beta)$.

On the other hand, $V_Q^{(2)}(\alpha)$ is a simple quotient of $V_Q^{(2)}(\gamma) \otimes V_Q^{(2)}(\beta)$ by Proposition 3.9. Hence we obtain the desired result $\mathcal{F}_Q^{(2)}(S_Q(\alpha)) \simeq V_Q^{(2)}(\alpha)$. □

Lemma 4.2. *Let* $\beta, \gamma \in \Delta^+_{\mathfrak{g}_0^{(1)}}$ *and* $t = 1, 2$. *If* $R^{\mathrm{norm}}_{V_Q^{(t)}(\beta), V_Q^{(t)}(\gamma)}(z)$ *has a pole at* $z = 1$, *then* $\gamma \prec_Q \beta$.

Proof. By Proposition 3.3, it is enough to prove the case $t = 1$. Set $\phi^{-1}(\beta, 0) = (i, a)$ and $\phi^{-1}(\gamma, 0) = (j, b)$. Assume that $R^{\mathrm{norm}}_{V_Q^{(t)}(\beta), V_Q^{(t)}(\gamma)}(z)$ has a pole at $z = 1$, or equivalently, the denominator $d_{ij}(z)$ has a zero at $z = (-q)^{b-a}$. Then we can easily see that $d(i, j) + 2 \le b - a$ by Theorem 2.2. Here $d(i, j)$ is the distance of i and j in the Dynkin diagram. Then $\gamma \preceq_Q \beta$ by the definition (2.6). □

For $\{m_\beta\}_{\beta \in \Delta^+_{\mathfrak{g}_0^{(1)}}} \in \left(\mathbb{Z}_{\ge 0}\right)^{\Delta^+_{\mathfrak{g}_0^{(1)}}}$, let us denote by $\overset{\to}{\bigotimes}_{\beta \in \Delta^+_{\mathfrak{g}_0^{(1)}}} (V_Q^{(2)}(\beta))^{\otimes m_\beta})$ the increasingly ordered tensor product with respect to a total order on $\Delta^+_{\mathfrak{g}_0^{(1)}}$ stronger than $\preceq_Q$. The isomorphism class of the module $\overset{\to}{\bigotimes}_{\beta \in \Delta^+_{\mathfrak{g}_0^{(1)}}} (V_Q^{(2)}(\beta))^{\otimes m_\beta}$ does not depend on the choice of such a total order by Lemma 4.2. Note also that $\overset{\to}{\bigotimes} (V_Q^{(2)}(\beta))^{\otimes m_\beta})$ has a simple head by Lemma 4.2 and [15, Theorem 2.2.1 (ii)].

Definition 4.3. The category $\mathscr{C}_Q^{(2)}$ is the full subcategory of $\mathscr{C}_{\mathfrak{g}^{(2)}}$ consisting of objects M such that any simple subquotient of M is isomorphic to $\mathrm{hd}(\overset{\to}{\bigotimes}_{\beta \in \Delta^+_{\mathfrak{g}_0^{(1)}}} (V_Q^{(2)}(\beta))^{\otimes m_\beta})$ for some $\{m_\beta\}_{\beta \in \Delta^+_{\mathfrak{g}_0^{(1)}}} \in \left(\mathbb{Z}_{\ge 0}\right)^{\Delta^+_{\mathfrak{g}_0^{(1)}}}$.

Note that there is a similar characterization of $\mathscr{C}_Q^{(1)}$.

By the definition, $\mathscr{C}_Q^{(2)}$ is stable under taking subquotients and extensions. Note that we have

(4.1) If $V^{(t)}(\varpi_i)_x$ belongs to $\mathscr{C}_Q^{(t)}$, then it is isomorphic to $V_Q^{(t)}(\beta)$ for some $\beta \in \Delta^+_{\mathfrak{g}_0^{(1)}}$.

Indeed, the dominant extremal weight of $\mathrm{hd}\big(\overset{\to}{\otimes}(V_Q^{(t)}(\beta))^{\otimes m_\beta}\big)$ is equal to $\sum_{\beta\in\Delta^+_{\mathfrak{g}_0^{(1)}}} m_\beta\varpi_{i_\beta}$ for $t=1$ (respectively, is equal to $\sum_{\beta\in\Delta^+_{\mathfrak{g}_0^{(1)}}} m_\beta\varpi_{i'_\beta}$ for $t=2$), where $\phi(i_\beta,p_\beta)=(\beta,0)$ and i'_β denotes the first component of $\pi_{\mathfrak{g}^{(1)}}(i_\beta,(-q)^{p_\beta})$ for $\beta\in\Delta^+_{\mathfrak{g}_0^{(1)}}$.

The proof of the following theorem works also for $\mathcal{F}_Q^{(1)}$, and it provides an alternative proof of [16, Corollary 4.3.6].

Theorem 4.4. *The functor $\mathcal{F}_Q^{(2)}$ sends a simple module to a simple module. Moreover, the functor $\mathcal{F}_Q^{(2)}$ induces a bijection between the set of simple modules in* $\mathrm{Rep}(R)$ *and the set of simple modules in $\mathscr{C}_Q^{(2)}$.*

Proof. By Theorem 2.5, every simple module M in $\mathrm{Rep}(R)$ is isomorphic to the image of the homomorphism

$$\mathbf{r}_{\underline{m}}\colon \overrightarrow{\nabla}_Q(\underline{m}) := \mathop{\circ}_{k=1}^{r} S_Q(\beta_k)^{\circ m_k} \to \overleftarrow{\nabla}_Q(\underline{m}) := \mathop{\circ}_{k=1}^{r} S_Q(\beta_{r-k+1})^{\circ m_{r-k+1}} \tag{4.2}$$

for a unique $\underline{m}\in\mathbb{Z}^r_{\ge 0}$. Moreover we have

$$[\overrightarrow{\nabla}_Q(\underline{m})] \in [\mathrm{Im}(\mathbf{r}_{\underline{m}})] + \sum_{\underline{m}'\prec_{\mathbb{Z}}\underline{m}} \mathbb{Z}_{\ge 0}[\mathrm{Im}(\mathbf{r}_{\underline{m}'})]. \tag{4.3}$$

Applying the functor $\mathcal{F}_Q^{(2)}$ to (4.2), we obtain

$$\overrightarrow{W}_Q(\underline{m}) := \mathop{\otimes}_{k=1}^{r} V_Q^{(2)}(\beta_k)^{\otimes m_k} \xrightarrow{\mathcal{F}_Q^{(t)}(\mathbf{r}_{\underline{m}})} \overleftarrow{W}_Q(\underline{m}) := \mathop{\otimes}_{k=1}^{r} V_Q^{(2)}(\beta_{r-k+1})^{\otimes m_{r-k+1}} .$$

Now we shall prove that

$$\mathrm{Im}\big(\mathcal{F}_Q^{(2)}(\mathbf{r}_{\underline{m}})\big) \simeq \mathcal{F}_Q^{(2)}\big(\mathrm{Im}(\mathbf{r}_{\underline{m}})\big) \text{ is simple and isomorphic to } \mathrm{hd}(\overrightarrow{W}_Q(\underline{m})). \tag{4.4}$$

If $\underline{m}$ is a minimal element of $\mathbb{Z}^r_{\ge 0}\setminus\{0\}$, i.e. $\underline{m}$ is a unit vector, then (4.4) follows from Theorem 4.1. Assume that $\underline{m}$ is not minimal. By Lemma 4.2, we can apply [15, Theorem 2.2.1 (ii)]. Then, $\overrightarrow{W}_Q(\underline{m})$ has a simple head which is equal to the image of any non-zero map from $\overrightarrow{W}_Q(\underline{m})$ to $\overleftarrow{W}_Q(\underline{m})$. Thus it is enough to show that $\mathcal{F}_Q^{(t)}(\mathbf{r}_{\underline{m}})$ is non-zero.

By the induction hypothesis on $\prec_{\mathbb{Z}}$, every composition factor of $\mathrm{Ker}\big(\mathcal{F}_Q^{(t)}(\mathbf{r}_{\underline{m}})\big)$ is of the form $\mathrm{hd}(\overrightarrow{W}_Q(\underline{m}'))$ for some $\underline{m}'\prec_{\mathbb{Z}}\underline{m}$. By [15, Theorem 2.2.1 (iii)], $\mathrm{hd}(\overrightarrow{W}_Q(\underline{m}))$ is isomorphic to $\mathrm{hd}(\overrightarrow{W}_Q(\underline{m}'))$ if and only if $\underline{m}'=\underline{m}$. Thus we conclude that $\mathrm{hd}(\overrightarrow{W}_Q(\underline{m}))$

cannot appear as a composition factor of $\mathrm{Ker}\big(\mathcal{F}_Q^{(t)}(\mathbf{r}_{\underline{m}})\big)$, which yields that $\mathcal{F}_Q^{(t)}(\mathbf{r}_{\underline{m}})$ is non-zero. $\square$

Corollary 4.5.

(i) *The functor* $\mathcal{F}_Q^{(2)}$ *induces a functor* $\mathrm{Rep}(R) \to \mathscr{C}_Q^{(2)}$.

(ii) *The functor* $\mathcal{F}_Q^{(2)}$ *is faithful.*

(iii) *The category* $\mathscr{C}_Q^{(2)}$ *is stable under taking tensor products.*

Proof. (i) follows immediately from Theorem 4.4.

(ii) Since $\mathcal{F}_Q^{(2)}$ sends simples to simples, the module $\mathrm{Im}(\mathcal{F}_Q^{(2)}(f))$ does not vanish for any non-zero homomorphism f. Hence $\mathcal{F}_Q^{(2)}(f)$ is non-zero.

(iii) follows from the fact that $\mathcal{F}_Q^{(2)}\colon \mathrm{Rep}(R) \to \mathscr{C}_Q^{(2)}$ sends a convolution product to a tensor product. $\square$

By this corollary, we have another characterization of $\mathscr{C}_Q^{(2)}$: $\mathscr{C}_Q^{(2)}$ is the smallest full subcategory of $\mathscr{C}_{\mathfrak{g}^{(2)}}$ such that it is stable under taking subquotients, extensions, tensor products and it contains $V_Q^{(2)}(\alpha_i)$ for all $i \in I_0^{(1)}$.

Corollary 4.6. *The functor* $\mathcal{F}_Q^{(2)}$ *induces a ring isomorphism*

$$\varphi^{(2)} : \mathrm{K}(\mathrm{Rep}(R)) \xrightarrow{\sim} \mathrm{K}(\mathscr{C}_Q^{(2)})$$

and hence a ring isomorphism

$$\varphi := \varphi^{(2)} \circ (\varphi^{(1)})^{-1} : \mathrm{K}(\mathscr{C}_Q^{(1)}) \xrightarrow{\sim} \mathrm{K}(\mathscr{C}_Q^{(2)}).$$

Moreover, the ring isomorphism φ *sends simples to simples and preserves the dimensions.*

Proof. The first assertion follows from the preceding corollary and the fact that $\mathcal{F}_Q^{(2)}$ sends a convolution product to a tensor product. For the last assertion on the dimension, we can apply the same induction argument of [18, Theorem 3.5.10] along with Proposition 3.5. $\square$

Conjecture 4.7. *The functor* $\mathcal{F}_Q^{(t)}\colon \mathrm{Rep}(R) \to \mathscr{C}_Q^{(t)}$ $(t = 1, 2)$ *is an equivalence of categories.*

5. PROOF OF THEOREM 3.10

In this section, we give a proof of Theorem 3.10 for each $\mathfrak{g}$, which we have postponed in Section 3.

Lemma 5.1. *Let* $\mathfrak{g}^{(1)} = A_N^{(1)}$ *or* $D_N^{(1)}$. *Assume that* (i, x), (j, y) *is a pair of elements of* $\mathscr{S}(\mathfrak{g}^{(1)})$ *such that they are connected by an arrow and that* $V^{(1)}(\varpi_i)_x$ *and* $V^{(1)}(\varpi_j)_y$ *are not dual to each other. Then there exist a Dynkin quiver of type* $\mathfrak{g}_0^{(1)}$, *a height function* ξ, $a \in \mathbf{k}^\times$ *and* (i, s), $(j, t) \in \Gamma_Q$ *such that* $((i, x), (j, y)) = ((i, a(-q)^s), (j, a(-q)^t))$.

Postponing the proof of this lemma, we shall prove Theorem 3.10.

Proof of Theorem 3.10. Assume that there is an epimorphism

$$V^{(1)}(i, x) \otimes V^{(1)}(j, y) \twoheadrightarrow V^{(1)}(k, z) \tag{5.1}$$

or an epimorphism

$$V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(i, x)) \otimes V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(j, y)) \twoheadrightarrow V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(k, z)). \tag{5.2}$$

Then (i, x) and $(j, y) \in \mathscr{S}(\mathfrak{g}^{(1)})$ are connected by an arrow. Moreover $V^{(1)}(i, x)$ and $V^{(1)}(j, y)$ are not dual to each other. Indeed, if $V^{(1)}(i, x)$ and $V^{(1)}(j, y)$ were dual to each other, then $V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(i, x))$ and $V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(j, y))$ would be also dual to each other by Proposition 3.3 (iii). Then, there would not exist an epimorphism (5.1) nor (5.2).

Hence, by Lemma 5.1, there exist a Dynkin quiver Q of type $\mathfrak{g}_0^{(1)}$, a height function ξ, $a \in \mathbf{k}^\times$ and (i, s), $(j, t) \in \Gamma_Q$ such that $((i, x), (j, y)) = ((i, a(-q)^s), (j, a(-q)^t))$. We may assume $a = 1$ without loss of generality. Then, $V^{(1)}(i, x)$ and $V^{(1)}(j, y)$ belong to $\mathscr{C}_Q^{(1)}$, and also $V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(i, x))$ and $V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(j, y))$ belong to $\mathscr{C}_Q^{(2)}$. Hence $V^{(1)}(k, z)$ belongs to $\mathscr{C}_Q^{(1)}$ or $V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(k, z))$ belongs to $\mathscr{C}_Q^{(2)}$. By (4.1), there are $\alpha, \beta, \gamma \in \Delta^+_{\mathfrak{g}_0^{(1)}}$ such that $V^{(1)}(i, x) = V_Q^{(1)}(\alpha)$, $V^{(1)}(j, y) = V_Q^{(1)}(\beta)$, $V^{(1)}(k, z) = V_Q^{(1)}(\gamma)$, $V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(i, x)) = V_Q^{(2)}(\alpha)$, $V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(j, y)) = V_Q^{(2)}(\beta)$ and $V^{(2)}(\pi_{\mathfrak{g}^{(1)}}(k, z)) = V_Q^{(2)}(\gamma)$. Let $f\colon S_Q(\alpha) \circ S_Q(\beta) \to S_Q(\beta) \circ S_Q(\alpha)$ be a non-zero morphism in $\mathrm{Rep}(R)$. Then $\mathrm{Im}(f)$ is the simple head of $S_Q(\alpha) \circ S_Q(\beta)$ and $\mathrm{Im}\big(\mathcal{F}_Q^{(t)}(f)\big)$ is the simple head of $V_Q^{(t)}(\alpha) \otimes V_Q^{(t)}(\beta)$. Therefore, for each $t = 1, 2$, we have the equivalence relations:

$$\begin{aligned}
&\text{there exists an epimorphism } V_Q^{(t)}(\alpha) \otimes V_Q^{(t)}(\beta) \twoheadrightarrow V_Q^{(t)}(\gamma) \\
&\Longleftrightarrow \mathrm{Im}\big(\mathcal{F}_Q^{(t)}(f)\big) \simeq V_Q^{(t)}(\gamma) \simeq \mathcal{F}^{(t)}\big(S_Q(\gamma)\big) \\
&\Longleftrightarrow \mathrm{Im}(f) \simeq S_Q(\gamma).
\end{aligned}$$

Thus we obtain Theorem 3.10. ☐

Before starting the proof of Lemma 5.1, let us remark on the Auslander-Reiten quiver Γ_Q.

The set of vertices $(\Gamma_Q)_0$ of Γ_Q is described as follows:

$$(\Gamma_Q)_0 = \{(i,p) \in I_0 \times \mathbb{Z} \mid \xi_i - 2m_i \le p \le \xi_i,\ p \equiv \xi_i \bmod 2\} \tag{5.3}$$
$$\text{where } m_i := \max\{k \in \mathbb{Z}_{\ge 0} \mid \tau^k(\gamma_i) \in \Delta^+_{\mathfrak{g}_0}\}.$$

Example 5.2.

(a) For a quiver $\overrightarrow{Q} := \underset{1}{\bullet} \rightarrow \underset{2}{\bullet} \rightarrow \bullet \cdots\cdots \rightarrow \underset{N-1}{\bullet} \rightarrow \underset{N}{\bullet}$ of type A_N, we have $m_i = N - i$ for all $i \in I_0$.

(b) For a quiver $\overleftarrow{Q} := \underset{1}{\bullet} \leftarrow \underset{2}{\bullet} \leftarrow \bullet \leftarrow\cdots\cdots \underset{N-1}{\bullet} \leftarrow \underset{N}{\bullet}$ of type A_N, we have $m_i = i - 1$ for all $i \in I_0$.

Note that, for any Dynkin quiver Q of type D_N and any $1 \le i \le N$, m_i's in (5.3) are given as follows ([27, Lemma 1.11]):

$$\begin{cases} m_i = N-2 & \text{if } 1 \le i \le N-2, \\ m_{N-1} = N-3,\ \ m_N = N-1 & \text{if } N \equiv 1 \bmod 2 \text{ and } \xi_N = \xi_{N-1}+2, \\ m_{N-1} = N-1,\ \ m_N = N-3 & \text{if } N \equiv 1 \bmod 2 \text{ and } \xi_{N-1} = \xi_N+2, \\ m_{N-1} = m_N = N-2 & \text{otherwise.} \end{cases} \tag{5.4}$$

Proof of Lemma 5.1. Without loss of generality, we may assume $(i,x), (j,y) \in \mathscr{S}_0(\mathfrak{g}^{(1)})$. Set $(i,x) = (i,(-q)^s)$ and $(j,y) = (j,(-q)^t)$. By the assumption, $(-q)^\ell$ is a root of $d_{i,j}(z)$ for $\ell = |s-t| > 0$. In this proof, we only consider the case when $i \ge j$ and $s < t$. The other cases can be proved by similar arguments.

(Case $\mathfrak{g} = A_N^{(1)}$) Note that the case $i + j = t - s = N + 1$ is excluded by the assumption. By (2.8), we have

$$i - j + 2 \le t - s \le \min(i+j, 2(N+1) - i - j) \text{ and } t - s \equiv i - j \bmod 2.$$

Hence, one of the following two conditions holds:

(i) $t - s \le i + j - 2$,

(ii) $t - s \le 2N - i - j$.

Indeed, otherwise, we have $i+j,\ 2N-i-j+2 \le t-s \le \min(i+j, 2(N+1)-i-j)$, which implies $i+j = t-s = N+1$. It is a contradiction.

In case (i), we set $\xi_k = t+k-j$. Then we have $m_k = k-1$ by Example 5.2 (b), and $t-i-j+2 = \xi_i - 2m_i \le s \le \xi_i = t+i-j$.

In case (ii), we set $\xi_k = t-k+j$. Then we have $m_k = N-k$ by Example 5.2 (a), and $t-2N+i+j = \xi_i - 2m_i \le s \le \xi_i = t-i+j$.

Now let us consider the case $\mathfrak{g} = D_N^{(1)}$.

(Case $\mathfrak{g} = D_N^{(1)}$, $1 \le j \le N-2$) In this case we may assume that $i \le N-1$. Choosing $\xi_{N-1} = \xi_N$, we may assume $m_k = N-2$. By (2.9), we have $i-j+2 \le t-s \le 2N-2+j-i$ and $t-s \equiv i-j \mod 2$. Hence, by taking $\xi_j = t$, and $\xi_i = t-i+j$, we have $\xi_i - 2m_i = t-i+j-2N+4 \le s \le \xi_i$ except the case $i=j$ and $t-s = 2N-2$. The last case is excluded by the assumption that $V(\varpi_i)_x$ and $V(\varpi_j)_y$ are not dual to each other.

(Case $\mathfrak{g} = D_N^{(1)}$, $i, j \in \{N-1, N\}$) By (2.9), we have $2 \le t-s \le 2N-2$ and $t \equiv s \mod 2$.

Assume first that N is even. Then the case $i-j = t-s-2(N-1) = 0$ is excluded by the assumption. By (5.4), we have $m_k = N-2$. Hence it is enough to take $\xi_j = t$ and $\xi_i = t - 2\delta(i \ne j)$. Then $\xi_i - 2m_i \le s \le \xi_i$.

Assume next that N is odd. Then the case $i \ne j$, $t-s = 2(N-1)$ is excluded. If $i = j$ then we take $m_i = N-1$, $\xi_i = t$. If $i \ne j$, then $0 \le t-s \le 2(N-2)$ and we take $\xi_i = \xi_j = t$, $m_i = N-2$. Then $\xi_i - 2m_i \le s \le \xi_i$. □

References

1. T. Akasaka and M. Kashiwara, *Finite-dimensional representations of quantum affine algebras*, Publ. RIMS. Kyoto Univ., **33** (1997), 839-867.
2. R. Bedard, *On commutation classes of reduced words in Weyl groups*, European J. Combin. **20** (1999), 483–505.
3. V. Chari, *Braid group actions and tensor products*, Int. Math. Res. Not. **2002**, no. 7, 357–382.
4. V. Chari and A. Pressley, *A Guide to Quantum Groups*. Cambridge University Press, 1994.
5. ______, *Yangians, integrable quantum systems and Dorey's rule*, Commun. Math. Phys. **181** (1996), no. 2, 265–302.
6. ______, *Quantum affine algebras and affine Hecke algebras*, Pacific J. Math. **174** (2) (1996), 295-326.
7. I. V. Cherednik, *A new interpretation of Gelfand-Tzetlin bases*, Duke Math. J. **54** (1987), 563-577.
8. P. Gabriel, *Auslander-Reiten sequences and Representation-finite algebras*, Representation theory, I (Proc. Workshop, Carleton Univ., Ottawa, Ont., 1979), pp. 1–71, Lecture Notes in Math., **831**, Springer, Berlin, 1980.

9. V. Ginzburg , N. Reshetikhin and E. Vasserot, *Quantum groups and flag varieties*, A.M.S. Contemp. Math. **175** (1994), 101-130.
10. D. Hernandez, *Kirillov-Reshetikhin conjecture: the general case*, Int. Math. Res. Not. **2010** no.7, 149–193.
11. D. Hernandez and B. Leclerc, *Cluster algebras and quantum affine algebras*, Duke Math. J. **154** (2010), 265.341.
12. ______, *Quantum Grothendieck rings and derived Hall algebras*, , to appear in J. reine angew. Math., arXiv:1109.0862 [math.QA].
13. M. Jimbo, *A q-analogue of* $U(\mathfrak{gl}_{N+1})$*, Hecke algebra, and the Yang-Baxter equation*, Lett. Math.Phys. **11** (1986), 247-252.
14. V. Kac, *Infinite Dimensional Lie algebras*, 3rd ed., Cambridge University Press, Cambridge, 1990.
15. S.-J. Kang, M. Kashiwara and M. Kim, *Symmetric quiver Hecke algebras and R-matrices of quantum affine algebras*, arXiv:1304.0323 [math.RT].
16. ______, *Symmetric quiver Hecke algebras and R-matrices of quantum affine algebras II*, arXiv:1308.0651 [math.RT], to appear in Duke Math. J.
17. S.-J. Kang, M. Kashiwara, M. Kim and S.-j. Oh, *Simplicity of heads and socles of tensor products*, arXiv:1404.4125 [math.RT], to appear in Compositio Math.
18. ______, *Symmetric quiver Hecke algebras and R-matrices of quantum affine algebras III*, arXiv:1406.0591 [math.RT].
19. ______, *Monoidal categorification of cluster algebras II*, arXiv:1502.06714.
20. M. Kashiwara, *On level zero representations of quantum affine algebras*, Duke. Math. J. **112** (2002), 117–175.
21. M. Khovanov and A. Lauda, *A diagrammatic approach to categorification of quantum groups I*, Represent. Theory **13** (2009), 309–347.
22. S. Kato, *PBW bases and KLR algebras*, arXiv:1203.5254v4 [math.QA].
23. M. Kim, *Khovanov-Lauda-Rouquier algebras and R-matrices*, Ph. D. thesis, Seoul National University, 2012.
24. P. McNamara, *Finite dimensional representations of Khovanov-Lauda-Rouquier algebras I: finite type*, to appear in J. reine angew. Math.; arXiv:1207.5860.
25. S.-j. Oh, *The Denominators of normalized R-matrices of types* $A^{(2)}_{2n-1}$, $A^{(2)}_{2n}$, $B^{(1)}_{n}$ *and* $D^{(2)}_{n+1}$, arXiv:1404.6715 [math.QA].
26. ______, *Auslander-Reiten quiver of type A and generalized quantum affine Schur-Weyl duality*, arXiv:1405.3336 [math.RT].
27. ______, *Auslander-Reiten quiver of type D and generalized quantum affine Schur-Weyl duality*, arXiv:1406.4555 [math.RT].
28. P. Papi, *A characterization of a special ordering in a root system*, Proc. Amer. Math. Soc. **120** (1994), no. 3, 661–665.
29. C. Ringel, *PBW-bases of quantum groups*, J. Reine Angew. Math. **470** (1996), 51–88.
30. R. Rouquier, *2-Kac-Moody algebras*, arXiv:0812.5023 [math.RT].

31. I. Schur, *Über Eine Klasse Von Matrizen, die Sich Einer Gegeben Matrix Zurodenen Lassen*, Ph.D. thesis (1901). Reprinted in Gesamelte Abhandlungen **1**, 1–70.
32. I. Schur, *Über die rationalen Darstellungen der allgemeinen linearen Gruppe*, Sitzungsberichte der Königlich Preussischen Adademie der Wissenschaften zu Berlin (1927), 58–75. Reprinted in Gessamelte Abhandlungen **3**, 68–85.
33. S. Zelikson, *Auslander-Reiten quivers and the Coxeter complex*, Algebr. Represent. Theory **8** (2005), no. 1, 35–55.

Department of Mathematical Sciences and Research Institute of Mathematics, Seoul National University, Gwanak-ro 1, Gwanak-gu, Seoul 151-747, Korea
E-mail address: `sjkang@snu.ac.kr`

Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, Japan, & Department of Mathematical Sciences and Research Institute of Mathematics, Seoul National University, Seoul 151-747, Korea
E-mail address: `masaki@kurims.kyoto-u.ac.jp`

School of Mathematics, Korea Institute for Advanced Study, Seoul 130-722, Korea
E-mail address: `mhkim@kias.re.kr`

Department of Mathematical Sciences, Seoul National University Gwanak-ro 1, Gwanak-gu, Seoul 151-747, Korea
E-mail address: `sejin092@gmail.com`